

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/260288429>

Evidence-based Interventions For Improvement Of Maternal And Child Nutrition: What Can Be Done And At What Cost?

ARTICLE *in* THE LANCET · JUNE 2013

Impact Factor: 45.22 · DOI: 10.1016/S0140-6736(13)60996-4

CITATIONS

177

READS

213

9 AUTHORS, INCLUDING:



[Arjumand Rizvi](#)

Aga Khan University, Pakistan

30 PUBLICATIONS 665 CITATIONS

SEE PROFILE



[Neff Walker](#)

Johns Hopkins University

145 PUBLICATIONS 8,653 CITATIONS

SEE PROFILE



[Patrick Webb](#)

Tufts University

130 PUBLICATIONS 2,463 CITATIONS

SEE PROFILE



[Robert E Black](#)

Johns Hopkins Bloomberg School of Public...

649 PUBLICATIONS 35,707 CITATIONS

SEE PROFILE

Maternal and Child Nutrition 2



Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost?

Zulfiqar A Bhutta, Jai K Das, Arjumand Rizvi, Michelle F Gaffey, Neff Walker, Susan Horton, Patrick Webb, Anna Lartey, Robert E Black, The Lancet Nutrition Interventions Review Group, and the Maternal and Child Nutrition Study Group

Maternal undernutrition contributes to 800 000 neonatal deaths annually through small for gestational age births; stunting, wasting, and micronutrient deficiencies are estimated to underlie nearly 3·1 million child deaths annually. Progress has been made with many interventions implemented at scale and the evidence for effectiveness of nutrition interventions and delivery strategies has grown since *The Lancet* Series on Maternal and Child Undernutrition in 2008. We did a comprehensive update of interventions to address undernutrition and micronutrient deficiencies in women and children and used standard methods to assess emerging new evidence for delivery platforms. We modelled the effect on lives saved and cost of these interventions in the 34 countries that have 90% of the world's children with stunted growth. We also examined the effect of various delivery platforms and delivery options using community health workers to engage poor populations and promote behaviour change, access and uptake of interventions. Our analysis suggests the current total of deaths in children younger than 5 years can be reduced by 15% if populations can access ten evidence-based nutrition interventions at 90% coverage. Additionally, access to and uptake of iodised salt can alleviate iodine deficiency and improve health outcomes. Accelerated gains are possible and about a fifth of the existing burden of stunting can be averted using these approaches, if access is improved in this way. The estimated total additional annual cost involved for scaling up access to these ten direct nutrition interventions in the 34 focus countries is Int\$9·6 billion per year. Continued investments in nutrition-specific interventions to avert maternal and child undernutrition and micronutrient deficiencies through community engagement and delivery strategies that can reach poor segments of the population at greatest risk can make a great difference. If this improved access is linked to nutrition-sensitive approaches—ie, women's empowerment, agriculture, food systems, education, employment, social protection, and safety nets—they can greatly accelerate progress in countries with the highest burden of maternal and child undernutrition and mortality.

Introduction

Stunting prevalence has been decreasing slowly and 165 million children were stunted in 2011.¹ Undernutrition, consisting of fetal growth restriction, stunting, wasting, and deficiencies of vitamin A and zinc, along with suboptimum breastfeeding, underlies nearly 3·1 million deaths of children younger than 5 years annually worldwide, representing about 45% of all deaths in this group.² Maternal and child obesity have also increased in many low-income and middle-income countries.³

In a comprehensive review of nutrition interventions, we previously assessed 43 nutrition-related interventions in detail and reported estimates of efficacy and effect for 11 core interventions.⁴ Much progress has been made since with many interventions implemented at scale, assessments of promising new interventions, and new delivery strategies. We used standard methods to do a comprehensive review of potential nutrition-specific interventions to address undernutrition and micronutrient deficiencies in women and children. We modelled the potential effect of delivery of these interventions on lives saved in the 34 countries with 90% of the global burden of stunted children, and estimated the effect of various delivery platforms that could enhance equitable scaling up of nutrition-specific interventions.

Selection of interventions for review

We selected several nutrition-specific interventions across the lifecycle for assessment of evidence of benefit (figure 1); these interventions included those affecting adolescents, women of reproductive age, pregnant women, newborn babies, infants, and children. We also reviewed the evidence for delivery platforms for nutrition interventions and other emerging interventions of interest for nutrition of women and children.

We identified and relied on the most recent reviews with good quality methods for all interventions and updated the evidence by incorporating newer studies, when available. For other identified interventions, when we did not find any relevant review, we did a de-novo review using the methodology described in panel 1.⁵ Additionally, we consulted the electronic library on nutrition actions (eLENA) for existing evidence used by WHO for development of guidelines and policies for action (appendix p 2).

Interventions to address adolescent health and nutrition

There is growing interest in adolescent health as an entry point to improve the health of women and children, especially because an estimated 10 million girls younger than 18 years are married each year.⁶ A range of

Published Online

June 6, 2013
[http://dx.doi.org/10.1016/S0140-6736\(13\)60996-4](http://dx.doi.org/10.1016/S0140-6736(13)60996-4)

This online publication has been corrected. The corrected version first appeared at thelancet.com on June 20, 2013

This is the second in a Series of four papers about maternal and child nutrition

Aga Khan University, Karachi, Pakistan (Prof Z A Bhutta PhD, J K Das MBA, A Rizvi MSc); Hospital for Sick Children, Toronto, ON, Canada (M F Gaffey MSc); Johns Hopkins University, Bloomberg School of Public Health, Baltimore, MD, USA (N Walker PhD, Prof R E Black PhD); University of Waterloo, Waterloo, ON, Canada (Prof S Horton PhD); Tufts University, Medford, MA, USA (Prof P Webb PhD); and University of Ghana, Accra, Ghana (Prof A Lartey PhD)

Correspondence to: Prof Zulfiqar A Bhutta, Center of Excellence in Women and Child Health, The Aga Khan University, Karachi 74800, Pakistan
zulfiqar.bhutta@aku.edu

See Online for appendix

Key messages

- Globally, 165 million children are stunted; undernutrition underlies 3·1 million deaths in children younger than 5 years.
- A clear need exists to introduce promising evidence-based interventions in the preconception period and in adolescents in countries with a high burden of undernutrition and young age at first pregnancies; however, targeting and reaching a sufficient number of those in need will be challenging.
- Promising interventions exist to improve maternal nutrition and reduce fetal growth restriction and small-for-gestational-age (SGA) births in appropriate settings in developing countries, if scaled up before and during pregnancy. These interventions include balanced energy protein, calcium, and multiple micronutrient supplementation and preventive strategies for malaria in pregnancy.
- Replacement of iron-folate with multiple micronutrient supplements in pregnancy might have additional benefits for reduction of SGA in at-risk populations, although further evidence from effectiveness assessments might be needed to guide a universal policy change.
- Strategies to promote breastfeeding in community and facility settings have shown promising benefits on enhancing exclusive breastfeeding rates; however, evidence for long-term benefits on nutritional and developmental outcomes is scarce.
- Evidence for the effectiveness of complementary feeding strategies is insufficient, with much the same benefits noted from dietary diversification and education and food supplementation in food secure populations and slightly greater effects in food insecure populations. Further effectiveness trials are needed in food insecure populations with standardised foods (pre-fortified or non-fortified), duration of intervention, outcome definition, and cost effectiveness.
- Treatment strategies for severe acute malnutrition with recommended packages of care and ready-to-use therapeutic foods are well established, but further evidence is needed for prevention and management strategies for moderate acute malnutrition in population settings, especially in infants younger than 6 months.
- Data for the effect of various nutritional interventions on neurodevelopmental outcomes are scarce; future studies should focus on these aspects with consistency in measurement and reporting of outcomes.
- Conditional cash transfers and related safety nets can address the removal of financial barriers and promotion of access of families to health care and appropriate foods and nutritional commodities. Assessments of the feasibility and effects of such approaches are urgently needed to address maternal and child nutrition in well supported health systems.
- Innovative delivery strategies, especially community-based delivery platforms, are promising for scaling up coverage of nutrition interventions and have the potential to reach poor populations through demand creation and household service delivery.
- Nearly 15% of deaths of children younger than 5 years can be reduced (ie, 1 million lives saved), if the ten core nutrition interventions we identified are scaled up.
- The maximum effect on lives saved is noted with management of acute malnutrition (435 000 [range 285 000–482 000] lives saved); 221 000 (135 000–293 000) lives would be saved with delivery of an infant and young child nutrition package, including breastfeeding promotion and promotion of complementary feeding; micronutrient supplementation could save 145 000 (30 000–216 000) lives.
- These interventions, if scaled up to 90% coverage, could reduce stunting by 20·3% (33·5 million fewer stunted children) and can reduce prevalence of severe wasting by 61·4%.
- The additional cost of achieving 90% coverage of these proposed interventions would be Int\$9·6 billion per year.

interventions exist in relation to adolescent health and nutrition, which could also affect the period before first pregnancy or between pregnancies. Evidence supporting reproductive health and family planning interventions in this age group suggests that it might be possible to reduce unwanted pregnancies and optimise age at first pregnancy. These aims might be important to reduce the risk of small-for-gestational age (SGA) births in populations in which a substantial proportion of births occur in adolescents. Opportunities might also exist to address micronutrient deficiencies and emerging issues of overweight and obesity in adolescents through community and school-based education platforms. Although evidence from robust randomised controlled trials is scarce, we identified a range of interventions in the adolescent period affecting maternal, newborn, and child health and nutrition outcomes (panel 2^{7–18}).

Interventions in women of reproductive age and during pregnancy

Folic acid supplementation

Neural tube defects can be effectively prevented with periconceptional folic acid supplementation. A review¹⁹ of five trials of periconceptional folic acid supplementation suggested a 72% reduction in risk of development of neural tube defects and a 68% reduction in risk of recurrence compared with either no intervention, placebo, or micronutrient intake without folic acid (table 1^{19–26}). A review²⁰ of folic acid supplementation during pregnancy showed that folic acid supplementation improved mean birthweight, with a 79% reduction in the incidence of megaloblastic anaemia (table 1^{19–26}). Furthermore no evidence of adverse effects was noted from folic acid supplementation in programme settings. Despite strong




Figure 1: Conceptual framework

WRA=women of reproductive age. WASH=water, sanitation, and hygiene. SAM=severe acute malnutrition. MAM=moderate AM.

evidence of benefit, reaching women of reproductive age in the periconceptual period to provide folic acid supplements through existing delivery platforms remains a logistical challenge. Fortification of cereals and other foods might be a feasible way to reach the population in need.

Iron or iron and folic acid supplementation

A review²¹ of iron supplementation in non-pregnant women of reproductive age showed that intermittent iron supplementation (alone or with any other vitamins and minerals) reduced the risk of anaemia by 27% (table 1^{19–26}). A Cochrane review²² of daily iron supplementation to women during pregnancy reported a 70% reduction in anaemia at term, a 67% reduction in iron deficiency anaemia (IDA), and 19% reduction in the incidence of low birthweight. Another review²⁷ further suggests that the effects were much the same in women receiving intermittent iron supplementation, or daily iron, or iron and folic acid supplementation. Although some evidence suggests that side-effects are fewer with intermittent iron therapy in non-anaemic populations, WHO recommends daily iron supplementation during pregnancy as part of the standard of care in populations at risk of iron deficiency.²⁸

Panel 1: Methods, search strategy, and selection criteria

As per the Child Health Epidemiology Reference Group (CHERG) systematic review guidelines,⁵ we searched PubMed, Cochrane libraries, electronic library on evidence on nutrition actions (eLENA), and WHO regional databases and included publications in every language available in these databases. We used Medical Subject Heading Terms (MeSH) and keyword search strategies with various combinations of relevant terms. We made every effort to gather unpublished data when reports were available for full abstraction. Inclusion and exclusion criteria were established for each area of review, and studies meeting these criteria were double data extracted and categorised according to outcome. Evidence was then summarised by outcome and study design, including study quality, generalisability, and summary outcome measures. We did meta-analyses for each outcome containing more than one study, using either the Mantel-Haenszel or the Der Simonian-Laird pooled relative risks (RR, with 95% CIs), when there was unexplained heterogeneity of effect. Heterogeneity was assessed by visual inspection of forest plots and by the χ^2 p value ($p < 0.10$). The binary measure for individual studies and pooled statistics was reported as the RR between the experimental and control groups with 95% CIs. For the outcome of interest for each intervention, we applied the CHERG Rules for Evidence Review⁵ to generate a final estimate.

Maternal multiple micronutrient supplementation

Multiple-micronutrient deficiencies often coexist in low-income and middle-income countries (LMICs) and can be exacerbated in pregnancy with potentially adverse maternal outcomes. A Cochrane review²³ of multiple

Panel 2: Interventions to address adolescent nutrition and preconception care

Women of lower socioeconomic status and young age are at risk of being undernourished and underweight. Ronnenberg and colleagues⁷ assessed the association between preconception anaemia and poor fetal and neonatal outcomes. They showed that the risk of being born low birthweight was significantly greater with moderate preconception anaemia (odds ratio [OR] 6.5, 95% CI 1.6–26.7) and fetal growth restriction (4.6, 1.5–13.5).

Important factors indirectly related to maternal, fetal, and neonatal nutritional status and pregnancy outcomes include young age at first pregnancy and repeated pregnancies. Young girls who are not physically mature might enter pregnancy with depleted nutrition reserves and anaemia.⁸ Adolescent pregnancy is associated with a 50% increased risk of stillbirths and neonatal deaths, and increased risk of preterm birth, low birthweight, and asphyxia.^{9–11} Adolescents are especially prone to complications of labour and delivery, such as obstructed and prolonged labour, vesico-vaginal fistulae, and infectious morbidity.¹¹ In societies in which most births are within a marital relationship, interventions to increase the age at marriage and first pregnancy are important.¹² Evidence suggests that programmes for adolescent mothers can reduce repeat adolescent pregnancies by 37% (95% CI 12–51%) when they teach parenting skills through home visitation and provide young mothers with education and vocational or job support.

Two reviews by Conde-Agudelo and colleagues^{13,14} assessed the association between inter-pregnancy intervals with maternal, newborn, and child health outcomes and found a J-shaped dose-response association for perinatal outcomes. Short inter-pregnancy intervals (<6 months) were associated with a higher probability of maternal anaemia (32%) and stillbirths (40%) whereas longer intervals (>60 months) were associated with an increased risk of pre-eclampsia.¹⁵ Both short and long birth intervals increase the risk for preterm births (OR 1.45 [95% CI 1.30–1.61] for short term; OR 1.21 [95% CI 1.12–1.30] for long intervals), low birthweight (OR 1.65 [95% CI 1.27–2.14] for short intervals; relative risk [RR] 1.37 [95% CI 1.21–1.55] for long intervals), and neonatal mortality (OR 1.31 [95% CI 0.96–1.79] for short interval; RR 1.15 [95% CI 1.06–1.25] for long intervals). With repeated pregnancies and advanced maternal age there is increased risk of chromosomal abnormalities, and increased risks of gestational diabetes and hypertension, stillbirths (RR 1.62, 95% CI 1.50–1.76), perinatal mortality (RR 1.44, 95% CI 1.10–1.89), and low birthweight (RR 1.61, 95% CI 1.16–2.24).^{16,17} These findings support the need to optimise age at first pregnancy and family size and inter-pregnancy intervals. A global unmet need exists for family planning with more than 100 million unmarried women in developing countries not using contraception.¹⁸ Optimisation of age at first pregnancy must be coupled with promotion of effective contraceptive use and exclusive breastfeeding, so that women can ideally space their pregnancies 18–24 months apart.

micronutrient supplementation in pregnant women assessed 23 trials and reported an 11–13% reduction in low birthweight and SGA births, whereas effects on anaemia and IDA were much the same when compared with iron and folic acid supplements (table 1). Despite earlier concerns about potential excess neonatal mortality with multiple micronutrient use,²⁹ present analyses suggest no adverse effects on maternal mortality, stillbirths, perinatal, and neonatal mortality with insufficient data for neuro-developmental outcomes. Although scarce, there are interesting data for benefits of maternal multiple micronutrient supplementation on growth in early childhood.³⁰ Preliminary data from a large trial³¹ comparing multiple micronutrient with iron-folate supplementation in

pregnancy in Bangladesh show a significant reduction in preterm births with no adverse effects. Inclusion of this study in our meta-analysis confirms the reduction in low birthweight (relative risk [RR] 0.88, 95% CI 0.85–0.91) and SGA (0.89, 0.83–0.96) and is also indicative of a small effect on preterm births (0.97, 0.94–0.99). These findings support the potential replacement of iron-folate supplements in pregnancy with multiple micronutrient supplements in populations at risk.

Maternal calcium supplementation

Gestational hypertensive disorders are the second leading cause of maternal morbidity and mortality and are associated with increased risk of preterm birth and fetal growth restriction.^{32,33} Calcium supplementation during pregnancy in women at risk of low calcium intake has been shown to reduce maternal hypertensive disorders and preterm birth. A Cochrane review by Hofmeyr and colleagues³⁴ assessed 13 trials and showed that calcium supplementation during pregnancy reduced the incidence of gestational hypertension by 35%, pre-eclampsia by 55%, and preterm births by 24% (table 1). These estimates have been updated in a review²⁴ of 15 randomised controlled trials, which also showed a 52% reduction in the incidence of pre-eclampsia and confirmed that these effects were only noted in populations at risk of low calcium intake.

Maternal iodine supplementation or fortification

In nearly all regions affected by iodine deficiency, use of iodised salt is the most cost-effective way to avert deficiency. A Cochrane review³⁵ suggests that although iodised salt is an effective means to improve iodine status, no conclusions can be drawn about physical and mental development in children and mortality. In some regions of the world with severe iodine deficiency, salt iodisation alone might not be sufficient for control of iodine deficiency in pregnancy; in these circumstances iodised oil supplementation during pregnancy can be a viable option (table 1). A review²⁵ of five randomised trials of iodised oil supplementation in pregnancy in iodine-deficient populations showed a 73% reduction in cretinism and a 10–20% increase in developmental scores in children. Existing evidence supports continued focus on effective universal salt iodisation for women of reproductive age and those who are pregnant. Further high-quality controlled studies are needed to address dosage and alternative strategies for iodine supplementation in different population groups and settings.

Addressing maternal wasting and food insecurity with balanced energy and protein supplementation

Maternal undernutrition is a risk factor for fetal growth restriction and adverse perinatal outcomes.¹ Several nutritional interventions have been assessed in such situations, including dietary advice to pregnant women, provision of balanced energy protein supplements, and

	Evidence reviewed	Setting	Estimates
Folic acid supplementation			
Women of reproductive age	Systematic review of five trials ¹⁹ of periconceptual folic acid supplementation	Developing and developed countries	Significant effects: NTDs (RR 0.28, 95% CI 0.15–0.52), recurrence of NTDs (RR 0.32, 95% CI 0.17–0.60) Non-significant effects: other congenital abnormalities, miscarriages, still births
Pregnant women	Systematic review of 31 trials ²⁰	Mostly developed countries	Significant effects: mean birthweight (MD 135.75, 95% CI 47.85–223.68), incidence of megaloblastic anaemia (RR 0.21, 95% CI 0.11–0.38) Non-significant effects: preterm birth, still births, mean predelivery haemoglobin, serum folate, red cell folate
Iron and Iron-folate supplementation			
Women of reproductive age	Systematic review of 21 RCTs and quasi-experimental studies ²¹	Developing and developed countries. Intervention mostly given in school settings. Mostly effectiveness studies	Intermittent iron supplementation Significant effects: anaemia (RR 0.73, 95% CI 0.56–0.95), serum haemoglobin concentration (MD 4.58 g/L, 95% CI 2.56–6.59), serum ferritin concentration (MD 8.32, 95% CI 4.97–11.66) Non-significant effects: iron deficiency, adverse events, depression
Pregnant women	Systematic review of 43 RCTs and quasi-experimental studies ²² (34 iron alone, eight iron-folate)	Developed and developing countries. Intervention delivered in community or at facility antenatal clinic. Mostly effectiveness studies	Daily iron-alone supplementation Significant effects: low birthweight (RR 0.81, 95% CI 0.68–0.97), birthweight (MD 30.81 g, 95% CI 5.94–55.68), serum haemoglobin concentration at term (MD 8.88 g/L, 95% CI 6.96–10.80), anaemia at term (RR 0.30, 95% CI 0.19–0.46), iron deficiency (RR 0.43, 95% CI 0.27–0.66), iron deficiency anaemia (RR 0.33, 95% CI 0.16–0.69), side-effects (RR 2.36, 95% CI 0.96–5.82) Non-significant effects: premature delivery, neonatal death, congenital anomalies Iron-folate supplementation Significant effects: birthweight (MD 57.7 g, 95% CI 7.66–107.79), anaemia at term (RR 0.34, 95% CI 0.21–0.54), serum haemoglobin concentration at term (MD 16.13 g/L, 95% CI 12.74–19.52) Non-significant effects: low birthweight, premature birth, neonatal death, congenital anomalies
MMN supplementation			
Pregnant women	Systematic review of 21 RCTs ²³	Developed and developing countries. Studies compared MMN with two or fewer micronutrients	Significant effects: low birthweight (RR 0.88, 95% CI 0.85–0.91), SGA (RR 0.89, 95% CI 0.83–0.96), preterm birth (RR 0.97, 95% CI 0.94–0.99) Non-significant effects: miscarriage, maternal mortality, perinatal mortality, stillbirths, and neonatal mortality Insufficient data for neurodevelopmental outcomes
Calcium supplementation			
Pregnant women	Systematic review of 15 RCTs ²⁴	Developed and developing countries. Mostly effectiveness trials	Significant effects: pre-eclampsia (RR 0.48, 95% CI 0.34–0.67), birthweight 85 g (95% CI 37–133), preterm birth (RR 0.76, 95% CI 0.60–0.97) Non-significant effects: perinatal mortality, low birthweight, neonatal mortality
Iodine through iodisation of salt			
Pregnant women	Systematic review of five RCTs ²⁵	Mostly developing countries. Mostly effectiveness trials	Significant effects: cretinism at 4 years of age (RR 0.27, 95% CI 0.12–0.60), developmental scores 10–20% higher in young children, birthweight 3.82–6.30% higher
Maternal supplementation with balanced energy protein			
Pregnant women	Systematic review of 16 RCTs and quasi-experimental studies ²⁶	Developing and developed countries	Significant effects: SGA (RR 0.66, 95% CI 0.49–0.89), stillbirths (RR 0.62, 95% CI 0.40–0.98), birthweight (MD 73g, 95% CI 30–117) Non-significant effects: Bayley mental scores at 1 year
NTD=neural tube defects. RR=relative risk. MD=mean difference. RCT=randomised controlled trial. MMN=multiple micronutrient. SGA=small-for-gestational age.			

Table 1: Review of nutrition interventions for women of reproductive age and during pregnancy

high protein or isocaloric protein supplementation. In other contexts, prescription and promotion of low energy diets to pregnant women who are either overweight or exhibit high weight gain in early gestation have been assessed.³⁶ Balanced energy protein supplementation, providing about 25% of the total energy supplement as protein, is deemed an important intervention for prevention of adverse perinatal outcomes in malnourished women.^{26,37} A Cochrane review³⁸ concluded that balanced energy protein supplementation reduced the incidence of SGA by 32% and risk of stillbirths by 45% (table 1). An updated meta-analysis showed that balanced energy protein supplementation increased birthweight by 73 g (95% CI 30–117) and reduced risk of SGA by 34%, with more pronounced effects in malnourished women.²⁶

Nutrition interventions in neonates

Delayed cord clamping

Early clamping of the umbilical cord after birth is a common practice and permits immediate transfer of the baby for care as required, whereas delaying of clamping allows continued blood flow between the placenta and the baby for a longer duration. A Cochrane review³⁹ suggested that delayed cord clamping in term neonates led to significant increase in newborn haemoglobin and higher serum ferritin concentration at 6 months of age (table 2^{39–45}). Another review⁴⁶ of studies in preterm neonates concluded that delayed cord clamping was associated with 39% reduction in need for blood transfusion and a lower risk of complications after birth. Although promising, these strategies have as yet not been assessed for effect or feasibility of implementation at scale in health systems.

	Evidence reviewed	Setting	Estimates
Delayed cord clamping			
Term neonates	Systematic review of 11 RCTs ³⁹	Developing and developed countries. 24 and 36 weeks' gestation at birth	Significant effects: increased newborn haemoglobin concentration (MD 2.17 g/dL, 95% CI 0.28–4.06) Non-significant effects: postpartum haemorrhage, severe postpartum haemorrhage Delayed cord clamping was associated with an increased requirement for phototherapy for jaundice
Preterm neonates	Systematic review of 15 RCTs ⁴⁰	Developing and developed countries	Significant effects: reduced need for blood transfusion (RR 0.61, 95% CI 0.46–0.81), decrease in intraventricular haemorrhage (RR 0.59, 95% CI 0.41–0.85), reduced risk of necrotising enterocolitis (RR 0.62, 95% CI 0.43–0.90) Peak bilirubin concentration was higher for delayed cord clamping group (MD 15.01 mmol/L, 95% CI 5.62–24.40)
Neonatal vitamin K administration			
Neonates	Systematic review of two RCTs for intramuscular vitamin K and 11 RCTs for oral vitamin K ⁴¹	Developing and developed countries	Significant effects: One dose of intramuscular vitamin K reduced clinical bleeding at 1–7 days and improved biochemical indices of coagulation status. Oral vitamin K also improved coagulation status
Vitamin A supplementation			
Very low birthweight infants	Systematic review of nine RCTs ⁴²	Developed countries	Significant effects: reduced number of deaths and oxygen requirement at 1 month of age. Non-significant effects: one large trial showed no significant effect on neurodevelopment assessment at 18–22 months of age
Term neonates	Systematic review of five RCTs and quasi-experimental studies ⁴³	Developing countries	Significant effects: reduction in infant mortality at 6 months (RR 0.86, 95% CI 0.77–0.97) Non-significant effects: infant mortality at 12 months (RR 1.03, 95% CI 0.87–1.23) Little data available for cause specific mortality, morbidity, vitamin A deficiency, anaemia, and adverse events
Kangaroo mother care for promotion of breastfeeding and care of preterm and SGA infants			
Healthy neonates	Systematic review of 34 RCTs ⁴⁴	Developing and developed countries	Significant effects: increase in breastfeeding at 1–4 months after birth (RR 1.27, 95% CI 1.06–1.53), increased breastfeeding duration (MD 42.55 days, 95% CI 1.69–86.79)
Preterm neonates	Systematic review of 16 RCTs ⁴⁵	Developing and developed countries	Significant effects: reduction in the risk of mortality (RR 0.60, 95% CI 0.39–0.93), reduction in nosocomial infection and sepsis (RR 0.42, 95% CI 0.24–0.73), reduction in hypothermia (RR 0.23, 95% CI 0.10–0.55), reduced length of hospital stay (MD 2.4 days, 95% CI 0.7–4.1)

RCT=randomised controlled trial. MD=mean difference. RR=relative risk. SGA=small-for-gestational age.

Table 2: Review of nutrition interventions in neonates

Neonatal vitamin K administration

Vitamin K deficiency can result in bleeding in the first weeks of life and vitamin K is commonly given prophylactically after birth for prevention of bleeding. In the absence of vitamin K prophylaxis there is a 0.4–1.7% risk of development of clinically significant bleeding. A Cochrane review⁴¹ suggested that one dose of intramuscular vitamin K, when compared with placebo, reduced clinical bleeding at 1–7 days of life, including bleeding after circumcision (table 2). Oral and intramuscular vitamin K had much the same effects on improved biochemical indices of coagulation status at 1–7 days. Currently, vitamin K administration after birth is largely restricted to births in health facilities; no information is available on the public health significance of vitamin K deficiency-related bleeding in LMICs or population-based programmes for prevention.

Neonatal vitamin A supplementation

A Cochrane review⁴² of oral or intramuscular vitamin A supplementation to very low birthweight infants showed reduced mortality and oxygen requirement at 1 month of age compared with placebo (table 2).⁴² Although neonatal vitamin A supplementation has also been shown to be effective in reduction of all-cause mortality by 6 months of age, evidence is conflicting, and might be related to maternal vitamin A status.⁴⁶ Although a Cochrane

review⁴³ did report a 14% reduction in the risk of infant mortality at 6 months of age, four more trials are currently underway in Asia and Africa, and researchers agree that these additional data will be needed before development of recommendations for neonatal vitamin A supplementation.

Kangaroo mother care

Kangaroo mother care denotes early skin-to-skin contact between mother and baby at birth or soon thereafter, plus early and continued breastfeeding, parental support, and early discharge from hospital. A Cochrane review⁴⁴ of 34 randomised controlled trials of early skin-to-skin care in healthy neonates showed a significant 27% increase in breastfeeding at 1–4 months of age and increased duration of breastfeeding (table 2). In a Cochrane review⁴⁵ of 16 randomised trials, kangaroo mother care in preterm neonates was associated with a 40% reduction in the risk of mortality, a 58% reduction in nosocomial infection or sepsis, and a 77% reduction in prevalence of hypothermia. The trials included in these analyses were done in health facilities; although kangaroo mother care might also be useful for home deliveries, there is not yet evidence of effectiveness in community settings. Kangaroo mother care was also shown to increase some measures of infant growth, breastfeeding, and mother-infant attachment,⁴⁵ but few studies provide objective evidence of any effect on early child development.

Nutrition interventions in infants and children

Promotion of breastfeeding and supportive strategies

WHO recommends initiation of breastfeeding within 1 h of birth, exclusive breastfeeding of infants till 6 months of age, and continued breastfeeding until 2 years of age or older.⁶⁷ However, global progress on this intervention is both uneven and suboptimum.⁴⁸ The exact scientific basis for the absolute early time window of feeding within the first hour after birth is weak.^{49,50} A systematic review⁵¹ suggests that breastfeeding initiation within 24 h of birth is associated with a 44–45% reduction in all-cause and infection-related neonatal mortality, and is thought to mainly operate through the effects of exclusive breastfeeding. We updated the previous review by Imdad and colleagues,⁵² which assessed the effect of promotion interventions on occurrence of breastfeeding, and concluded that counselling or educational interventions increased exclusive breastfeeding by 43% at day 1, by 30% till 1 month, and by 90% from 1–5 months. Significant reductions in occurrence of mothers not breastfeeding were also noted; 32% reduction at day 1, 30% till 1 month, and 18% for 1–5 months⁵³ (table 3^{53–62}). Combined individual and group counselling seemed to be better than individual or group counselling alone. Although these results show the potential for scaling up, none of these trials address the issues of barriers around work environments and supportive strategies such as maternity leave provision. A Cochrane review⁶³ of interventions in the workplace to support breastfeeding for women found no trials. Although some trials are underway, much more needs to be done to assess innovations and strategies to promote breastfeeding in working women, especially in underprivileged communities.

Promotion of dietary diversity and complementary feeding

Complementary feeding for infants refers to the timely introduction of safe and nutritionally rich foods in addition to breast-feeding at about 6 months of age and typically provided from 6 to 23 months of age.⁶⁴ Different approaches have been used to create indicators of dietary diversity and to study its association with child malnutrition. In seven Latin American surveys, Ruel and Menon⁶⁵ noted significant associations between complementary feeding practices and height-for-age Z scores (HAZ). Similarly, analysis of Demographic Health Survey data to create a dietary diversity score based on seven food groups showed that increased dietary diversity was positively associated with height-for-age HAZ in nine of 11 countries.⁶⁶ More recently, WHO infant and young child feeding indicators were studied in 14 Demographic Health Survey datasets from low-income countries;⁶⁷ consumption of a minimum acceptable diet with dietary diversity reduced the risk of both stunting and underweight whereas minimum meal frequency was associated with lower risk of underweight only.

In an update of a previous review of complementary feeding,⁶⁸ we assessed 16 randomised and non-randomised controlled trials and programmes of moderate quality (table 3).⁵⁴ We identified ten studies that assessed the effect of nutrition education and seven studies that assessed the effect of provision of additional complementary foods (one trial with three intervention groups was in both these categories). Studies of nutrition education in food secure populations showed a significant increase in height (standard mean difference [SMD] 0·35, 95% CI 0·08–0·62, four studies), and HAZ (0·22, 0·01–0·43, four studies), whereas the effect on stunting was not statistically significant (RR 0·70, 95% CI 0·49–1·01, four studies). We identified a significant effect on weight gain (SMD 0·40, 95% CI 0·02–0·78, four studies), whereas no effects were noted for weight-for-age Z scores (WAZ; 0·12; 95% CI –0·02 to 0·26, four studies). Studies of nutrition education in food insecure populations (with an average daily per person income of less than US\$1·25) showed significant effects on HAZ (SMD 0·25, 95% CI 0·09–0·42, one study), stunting (RR 0·68, 95% CI 0·60–0·76, one study), and WAZ (SMD 0·26, 95% CI 0·12–0·41, two studies). The review⁵⁴ did not find any eligible study that provided complementary feeding (with or without education) in a food secure population. Overall, the provision of complementary foods in food insecure populations was associated with significant gains in HAZ (SMD 0·39; 95% CI 0·05–0·73, seven studies) and WAZ (SMD 0·26, 95% CI 0·04–0·48, three studies), whereas the effect on stunting did not reach statistical significance (RR 0·33, 95% CI 0·11–1·00, seven studies).

Vitamin A supplementation in children

A Cochrane review⁵⁵ of 43 randomised trials showed that vitamin A supplementation reduced all-cause mortality by 24% and diarrhoea-related mortality by 28% in children aged 6–59 months (table 3). Vitamin A supplementation also reduced the incidence of diarrhoea and measles in this age group but there was no effect on mortality and morbidity related to respiratory infections. Although a large effectiveness study⁶⁹ from India assessing the effect of vitamin A supplementation and deworming over several years did not show a significant effect on mortality from vitamin A supplementation (mortality ratio 0·96, 95% CI 0·89–1·03), inclusion of these data with previous results still shows a significant, albeit lower, effect on mortality (RR 0·88, 95% CI 0·84–0·94).⁵⁵ We believe that vitamin A supplementation continues to be an effective intervention in children aged 6–59 months in populations at risk of vitamin A deficiency.

Iron supplementation in infants and children

A Cochrane review⁵⁶ of 33 studies showed that intermittent iron supplementation to children younger than 2 years reduced the risk of anaemia by 49% and iron

	Setting	Estimates
Breast feeding promotion in infants		
Systematic review of 110 RCTs and quasi-experimental studies ⁵³	Developing and developed countries	Significant effects: educational or counselling interventions increased EBF by 43% (95% CI 9–87) at day 1, by 30% (19–42) till 1 month, and by 90% (54–134) from 1–6 months. Significant reductions in rates of no breastfeeding also noted; 32% (13–46) at day 1, 30% (20–38) 0–1 month, and 18% (11–23) for 1–6 months. Non-significant effects: predominant and partial breastfeeding
Complementary feeding promotion in children 6–24 months of age		
16 RCTs and quasi-experimental studies ⁵⁴	Mostly from food secure populations. Various foods used	Nutrition education in food secure populations Significant effects: increased height gain (SMD 0.35; 95% CI 0.08–0.62), HAZ (SMD 0.22; 95% CI 0.01–0.43), weight gain (SMD 0.40, 95% CI 0.02–0.78) Non-significant effects: stunting, WAZ Nutrition education in food insecure populations Significant effects: HAZ (SMD 0.25, 95% CI 0.09–0.42), stunting (RR 0.68, 95% CI 0.60–0.76), WAZ (SMD 0.26, 95% CI 0.12–0.41) Complementary food provision with or without education in food insecure populations Significant effects: HAZ (SMD 0.39, 95% CI 0.05–0.73), WAZ (SMD 0.26, 95% CI 0.04–0.48) Non-significant effects: stunting (RR 0.33, 95% CI 0.11–1.00)
Preventive vitamin A supplementation in children 6 months to 5 years of age		
Systematic review of 43 RCTs ⁵⁵	Developing and developed countries	Significant effects: reduced all-cause mortality (RR 0.76, 95% CI 0.69–0.83), reduced diarrhoea-related mortality (RR 0.72, 95% CI 0.57–0.91), reduced incidence of diarrhoea (RR 0.85, 95% CI 0.82–0.87), reduced incidence of measles (RR 0.50, 95% CI 0.37–0.67) Non-significant effects: measles-related and ARI-related mortality
Iron supplementation in children		
Systematic review of 33 RCTs and quasi-experimental studies ⁵⁶	LMICs. Participant's ages ranged from neonates to 19 years	Intermittent iron supplementation Significant effects: decreased anaemia (RR 0.51, 95% CI 0.37–0.72), decreased iron deficiency (RR 0.24, 95% CI 0.06–0.91), increased haemoglobin concentration (MD 5.20 g/L, 95% CI 2.51–7.88), increased ferritin concentration (MD 14.17 mcg/L, 95% CI 3.53–24.81) Non-significant effects: HAZ, WAZ Evidence for mental development, motor skill development, school performance, and physical capacity was assessed by very few studies and showed no clear effect
Systematic review of 17 RCTs ⁵⁷	Developing and developed countries. In children aged 6 months to 15 years	Significant effects: increased mental development score (SMD 0.30, 95% CI 0.15–0.46), increased intelligence quotient scores (≥ 8 years age; SMD 0.41, 95% CI 0.20–0.62) Non-significant effects: Bayley mental development index in younger children (≤ 27 months old), motor development
MMN supplementation including iron in children		
Systematic review of 18 trials ⁵⁸	Mostly developing countries. In children aged 6 months to 16 years	MMN supplementation Significant effects: increased length (MD 0.13, 95% CI 0.06–0.21), increased weight (MD 0.14, 95% CI 0.03–0.25) MMN might be associated with marginal increase in fluid intelligence and academic performance in healthy school children
Systematic review of 17 RCTs ⁵⁹	Developing countries. Mostly effectiveness studies. In children aged 6 months to 11 years	Micronutrient powders Significant effects: Reduced anaemia (RR 0.66, 95% CI 0.57–0.77), reduced iron deficiency anaemia (RR 0.43, 95% CI 0.35–0.52), reduced retinol deficiency (RR 0.79, 95% CI 0.64–0.98). Improved haemoglobin concentrations (SMD 0.98, 95% CI 0.55–1.40). MNP was associated with a significant increase in diarrhoea (RR 1.04, 95% CI 1.01–1.06) Non-significant effects: serum ferritin, zinc deficiency, stunting, wasting, underweight, HAZ, WAZ, WHZ, fever, URI
Zinc supplementation in children		
Systematic review of 18 RCTs ^{60,61}	Mostly developing countries. In children younger than 5 years	Preventive zinc supplementation Significant effects: mean height improved by 0.37 cm (SD 0.25) in children supplemented for 24 weeks, diarrhoea reduced by 13% (95% CI 6–19), pneumonia reduced by 19% (95% CI 10–27) Non-significant effects: mortality (cause specific and all-cause)
Systematic review of 13 trials ⁶²	Developing and developed countries. In children younger than 5 years	Non-significant effects: Mental developmental index, psychomotor development index

RCT=randomised controlled trial. EBF=exclusive breastfeeding. HAZ=height-for-age Z score. WAZ=weight-for-age Z score. WHZ=weight-for-height Z score. MMN=multiple micronutrient. ARI=acute respiratory infection. URI=upper-respiratory infection. SMD=standard mean difference. MD=mean difference. RR=relative risk.

Table 3: Review of evidence for nutrition interventions for infants and children

deficiency by 76% (table 3). The findings also suggested that intermittent iron supplementation could be a viable public health intervention in settings in which daily supplementation had not been implemented or was not feasible. A review⁵⁷ of the effect of iron supplementation in children on mental and motor development showed only small gains in mental development and intelligence scores in supplemented school-age children who were initially anaemic or iron-deficient. There was no convincing evidence that iron

treatment had an effect on mental development in children younger than 27 months.

Since the demonstration of increased risk of admission to hospital and serious illnesses with iron supplementation,⁷⁰ there has been concern about administration of iron supplements in malaria endemic areas. WHO currently recommends administration of iron supplements in malaria endemic areas on the stipulation that malaria prevention and treatment is made available.⁷¹

Multiple micronutrient supplementation in children

Although the theoretical benefits of strategies to improve diet quality and micronutrient density of foods consumed by small children are well recognised, few resource-poor countries have clear policies in support of integrated strategies to control micronutrient deficiencies in young children.⁷² Available options include the provision of multiple micronutrients via supplements, micronutrient powders, or fortified ready-to-use foods including lipid-based nutrient supplements. A comprehensive review of the effects of multiple micronutrients compared with two or fewer micronutrients showed small benefits on linear growth (mean difference [MD] 0.13, 95% CI 0.06–0.21) and weight gain (0.14, 0.03–0.25) but with little evidence of effect on morbidity outcomes as suggested by individual studies (table 3).⁵⁸ Another review⁷³ of the effect of multiple micronutrient supplementation on improvement of cognitive performance in children concluded that multiple micronutrient supplementation might be associated with a marginal increase in reasoning abilities but not with acquired skills and knowledge.

Micronutrient powders are increasingly in use at scale in programmes to address iron and multiple micronutrient deficiencies in children. We reviewed 16 randomised controlled trials to assess the effectiveness of micronutrient powders and estimated that they significantly improved haemoglobin concentration and reduced IDA by 57% and retinol deficiency by 21%.⁵⁹ We noted no evidence of benefit on linear growth. However, in-line with findings from an earlier review of liquid iron supplementation trials,⁷⁴ use of micronutrient powders was shown to be associated with a significant increase in the incidence of diarrhoea (RR 1.04, 95% CI 1.01–1.06), largely because of results from a recent large cluster-randomised controlled trial of micronutrient powders in Pakistan in malnourished children.⁷⁵ These findings underscore the need for further assessment of micronutrient powder programmes in varying contexts for safety and benefits.

Preventive zinc supplementation in children

Preventive zinc supplementation in populations at risk of zinc deficiency reduces the risk of morbidity from childhood diarrhoea and acute lower respiratory infections and might increase linear growth and weight gain in infants and young children.^{60,76} A review by Yakoob and colleagues⁶¹ assessed 18 studies from developing countries and showed that preventive zinc supplementation reduced the incidence of diarrhoea by 13% and pneumonia by 19%, with a non-significant 9% reduction in all-cause mortality (table 3). However, subgroup analysis showed that there was a significant 18% reduction in all-cause mortality in children aged 12–59 months. A daily dose of 10 mg zinc per day over 24 weeks in children younger than 5 years could lead to an estimated net gain of 0.37 cm (SD 0.25) in height in

zinc-supplemented children compared with placebo.⁶⁰ There is no convincing evidence that zinc supplementation in infants or children results in improved motor or mental development.⁶²

Disease prevention and management

Several interventions have the potential to affect health and nutrition outcomes through reduction in the burden of infectious diseases. Table 4^{77–88} summarises the evidence for interventions for disease prevention and management.

Prevention and treatment of severe acute malnutrition

A substantial global burden of wasting exists, especially severe acute malnutrition (SAM; weight-for-height Z score [WHZ] <–3), which coexists with moderate acute malnutrition (MAM; WHZ <–2). In stable non-emergency situations with endemic malnutrition, MAM can often present in combination with stunting. Most of the interventions previously discussed should be implemented to prevent the development of SAM in food insecure populations. Several approaches for prevention and treatment are in use. Although the provision of complementary and supplementary foods could be considered in targeted food distribution programmes, other ways to stimulate access and purchasing power can be conceived. Where markets are fragmented or food access is constrained, appropriate food supplements might be considered as in-kind transfers. WHO recommends inpatient treatment for children with complicated SAM, with stabilisation and appropriate treatment of infections, fluid management, and dietary therapy and also supports community-based care for uncomplicated SAM.⁸⁹ Although facility-based treatment of SAM remains important, community management of SAM continues to grow rapidly globally. This shift in treatment norms from centralised, inpatient care towards community-based models allows more affected children to be reached and is cost effective. Up to an estimated 15% of cases of SAM will need initial facility-based care, whereas the rest can receive only community-based treatment.⁹⁰

Facility-based management of SAM according to the WHO protocol

A scientific literature review by Schofield and Ashworth⁹¹ showed that between the 1950s and 1990s, case fatality rates were typically 20–30% in children with SAM treated in hospitals or rehabilitation units, and rates were higher (50–60%) for oedematous malnutrition. A previous review⁴ of existing studies had estimated that following the WHO protocol, as opposed to standard care, would lead to a 55% reduction in deaths (RR 0.45, 95% CI 0.32–0.62; random effects).

In view of the limitations of analysis and variable quality of studies in the previous review, we updated the review to assess the effect of the WHO protocol or adaptations thereof on recovery and case fatality of children with SAM.

Case fatality rates ranged from 3·4% to 35%. The highest case fatality rate stemmed from a cohort of children with HIV infection.^{92,93} Only two studies provided information on recovery rates, which were 79·7% and 83·3%.^{94,95} In summary, the WHO protocol is substantiated through much evidence, based both on research and expert opinion. However, a clear need exists for continued work to improve staff training and quality⁹⁶ to achieve high rates of survival across various resource-constrained settings.

Community-based management of SAM

The products used to deliver nutrients for management of SAM and MAM, and the approaches used to target and deliver these products, evolved rapidly during the past decade. Innovations include new formulations and packaging and a shift from institutional to community-based management.

We reviewed interventions to treat SAM in community settings, and were largely able to pool studies comparing

	Settings	Estimates
WASH interventions		
Overview of three systematic reviews ⁷⁷	Developing countries	Significant effects: reduced risk of diarrhoea with hand washing with soap (RR 0·52, 95% CI 0·34–0·65), with improved water quality, and with excreta disposal
DHS data from 65 countries ⁷⁸	Developing countries	Significant effects: a recent World Bank report ⁷⁸ based on analysis of trends in DHS data suggests that open defecation explained 54% of international variation in child height by contrast with GDP, which only explained 29%. A 20 percentage point reduction in open defecation was associated with a 0·1 SD increase in child height A Cochrane review of the effect of WASH interventions on nutrition outcomes is underway ⁸⁷
Maternal deworming		
Systematic review of five RCTs ⁷⁹	Developing countries	Non-significant effects: one dose of anthelmintic in second trimester of pregnancy had a non-significant effect on maternal anaemia, low birthweight, preterm births, and perinatal mortality
Deworming in children (for soil-transmitted intestinal worms)		
Systematic review of 34 RCTs ⁸⁰	Developing countries	Non-significant effects: one-dose deworming had a non-significant effect on haemoglobin and weight gain. For multiple doses at 1 year follow up, there was a non-significant effect on weight, haemoglobin, cognition, and school attendance Treatment after confirmed infection Significant effects: one-dose of deworming drugs increased weight (0·58 kg, 95% CI 0·40–0·76) and haemoglobin (0·37 g/dL, 95% CI 0·1–0·64). Evidence on cognition was inconclusive These analyses are corroborated by the large-scale DEVTA trial ⁸⁸ of regular deworming and VAS over 5 years, which also did not show any benefits on weight gain or mortality
Feeding practices in diarrhoea		
Review of 29 RCTs ⁸¹	Developing countries	Significant effects: in acute diarrhoea, lactose-free diets, when compared with lactose-containing diets, significantly reduced incidence of diarrhoea (SMD –0·36, 95% CI –0·62 to –0·10) and treatment failure (RR 0·53, 95% CI 0·40–0·70) Non-significant effects: weight gain
Zinc therapy for diarrhoea		
Systematic review of 13 studies ⁸²	Mostly Asia	Significant effects: reduced all-cause mortality reduced by 46% (95% CI 12–68), diarrhoea-related admissions to hospital by 23% (95% CI 15–31) Non-significant effects: diarrhoea-specific mortality, diarrhoea-prevalence Zinc reduced duration of acute diarrhoea by 0·50 days and persistent diarrhoea by 0·68 days
IPTp/ITN for malaria in pregnancy		
Systematic review of 16 RCTs ⁸³	Mostly Africa	Significant effects: Anti-malarials to prevent malaria in all pregnant women reduced antenatal parasitemia (RR 0·53, 95% CI 0·33–0·86), increased birthweight (MD 126·7 g, 95% CI 88·64–164·75), reduced low birthweight by 43% (RR 0·57, 95% CI 0·46–0·72) and severe antenatal anaemia 38% (RR 0·62, 95% CI 0·50–0·78) Non-significant effects: perinatal deaths
Systematic review of six RCTs ⁸⁴	Developing countries	Significant effects: ITNs in pregnancy reduced low birthweight (RR 0·77, 95% CI 0·61–0·98) and reduced fetal loss (first to fourth pregnancy; RR 0·67, 95% CI 0·47–0·97) Non-significant effects: anaemia and clinical malaria
Malaria prophylaxis in children		
Systematic review of seven RCTs ⁸⁵	Developing countries of West Africa	Significant effects: Reduced clinical malaria episodes (RR 0·26; 95% CI 0·17–0·38), reduced severe malaria episodes (RR 0·27, 95% CI 0·10–0·76). IPTc also reduced risk of moderately severe anaemia (RR 0·71, 95% CI 0·52–0·98) Non-significant effects: all-cause mortality
Systematic review of 22 RCTs ⁸⁶	Developing countries in Africa	Significant effects: ITNs improved packed cell volume of children by 1·7 absolute packed cell volume percent. When the control group used untreated nets, the difference was 0·4 absolute packed cell volume percent. ITNs and IRS reduced malaria-attributable mortality in children (1–59 months) by 55% (95% CI 49–61) in <i>Plasmodium falciparum</i> settings
WASH=water, sanitation, and hygiene. RCT=randomised controlled trial. DHS=Demographic and Health Survey. GDP=gross domestic product. RR=relative risk. MD=mean difference. SMD=standard mean difference. WAZ=weight-for-age Z score. HAZ=height-for-age Z score. DEVTA=de-worming and enhanced vitamin A. IPTp=intermittent preventive treatment of malaria in pregnancy. IPTc=IPT in children. ITN=insecticide-treated bednets. IRS=indoor residual spraying.		
Table 4: Review of evidence for disease prevention and management		

ready-to-use therapeutic foods (RUTF) with standard care, as opposed to rigorous evaluation of effectiveness of the approach in programme settings.⁹⁷ We identified no significant differences in mortality; however, children who received RUTF had faster rates of weight gain and had 51% greater likelihood to recover (defined as attaining WHZ ≥ -2) than did those receiving standard care.

Notably, a new randomised controlled trial⁹⁸ compared standard RUTF with RUTF and additional 7 day course of antibiotics, either amoxicillin or cefdinir, in children with uncomplicated SAM. This trial showed that the children receiving an antibiotic had a lower mortality rate, faster recovery rate, and higher weight gain compared with children receiving placebo. Although further research on this topic is needed, especially in children with HIV infection, this study shows that effective community management of SAM might require an approach that goes beyond merely the choice of specially formulated foods to the entire package of care.

Substantial programmatic evidence supports use of RUTF for community-based treatment,⁹⁹ which has substantially changed the approach to treatment of SAM. Yet because of the nature of the evidence, establishing effect estimates for the overall approach to community management has proved challenging. Available evidence shows some positive effects with the use of RUTF compared with standard care for the treatment of SAM in community settings, yet the differences were for the most part small and several outcomes had substantial heterogeneity. An emphasis not only on the choice of commodities, but also on the quality of programme design and implementation is crucial to improvement of outcomes for children with SAM, as is research to fill information gaps, such as optimum treatment methods and approaches for treatment of breastfed infants younger than 6 months.

Interventions for prevention and management of obesity

Obesity is increasing in many populations and is one of the most important challenges of the 21st century. Obese women are at an increased risk of adverse pregnancy outcomes. A Cochrane review¹⁰⁰ assessed the effectiveness of interventions (eating, exercise, behaviour modification, or counselling) that reduce weight in obese pregnant women and identified no evaluable trials. Some studies assessed the effect of diet, exercise, or both for weight reduction in women after childbirth, and showed that women who exercised did not lose significantly more weight, but women who took part in a diet (MD -1.70 kg, 95% CI -2.08 to -1.32), or diet plus exercise programme (-2.89 kg; -4.83 to -0.95), did so. These interventions did not seem to adversely affect breastfeeding performance in any setting.¹⁰¹

We identified six reviews that examined breastfeeding in infancy and its association with obesity prevalence or average body-mass index (BMI) in childhood or

adulthood.^{102–107} All studies suggested a small protective effect of breastfeeding on obesity later in life, although the magnitude of the effect varied between reviews and the strength of the affect of confounding was unclear. The largest prospective follow up study in healthy term infants in Belarus showed that improving the duration and exclusivity of breastfeeding did not prevent overweight or obesity in children, nor did it affect insulin-like growth factor I concentrations at age 11.5 years.¹⁰⁸ These findings suggest that despite the myriad advantages of breastfeeding, population strategies to increase the duration and exclusivity of breastfeeding are unlikely to curb the obesity epidemic.

A Cochrane review¹⁰⁷ examined the effects of obesity prevention interventions delivered for more than 12 weeks on changes in BMI and BMI Z scores in children and suggested a significant beneficial effect across age groups with a SMD of -0.15 kg/m (95% CI -0.21 to -0.09). The subgroup analysis showed significant effects for children aged 6–12 years with non-significant effects in younger children and adolescents. Interventions that combined physical activity and diet were more effective than either delivered alone. Findings suggested that short-term interventions (<12 months duration) were more effective than were those delivered over a longer duration (SMD -0.17 , 95% CI -0.25 to -0.09 and SMD -0.12 , 95% CI -0.21 to -0.03 , respectively); however, there was substantial heterogeneity in all pooled estimates. Another review¹⁰⁹ of interventions to treat obesity in children showed that combined behavioural and lifestyle interventions or self-help could benefit overweight children and adolescents. Overall the evidence of effectiveness of all obesity prevention and therapeutic interventions is weak, underscoring the need for high-quality research in this discipline.

Delivery platforms and strategies for implementation of nutrition-specific interventions

Delivery strategies are crucial to achieve coverage with nutrition-specific interventions and to reach populations in need. A range of channels can provide opportunities for scaling up and reaching large segments of the population.

Fortification of staple foods and specific foods

A detailed discussion of fortification strategies is beyond the scope of this review. As supported by the Copenhagen consensus,¹¹⁰ fortification is one the most cost-effective strategies to reach populations at large. Further discussion of fortification as a means for delivery of key micronutrients is provided in panel 3^{111–118} and the accompanying report by Stuart Gillespie and colleagues.¹¹⁹

Cash transfer programmes

Financial incentives are widely used as policy strategies to ameliorate poverty, reduce financial barriers, and improve

Panel 3: Effect of fortification strategies

Food fortification is safe and cost effective in the prevention of micronutrient deficiencies and has been widely practised in developed countries for more than a century.¹¹¹ Foods can be fortified at three levels: mass or universal, targeted, and household. Mass or universal fortification—ideally legislated to be mandatory for industries—has the potential to produce foods and food products that are widely consumed by the general population (eg, salt iodisation and flour fortification with iron and folate). Mass fortification is by far the most cost-effective nutrition intervention, particularly when produced by medium-to-large scale industries.¹¹¹ Targeted fortification (eg, nutrient-fortified complementary foods for children aged 6–24 months) is important for subgroups of nutritionally vulnerable populations and populations in emergency situations whose nutrient intake is insufficient through available diets. Targeted fortification is also effective in resource poor settings where family foods do not include animal-source foods that are typically necessary to meet nutrient requirements of young children. Home fortification involves addition of nutrients directly to food consumed by women or children, or both, in the form of micronutrient powders or small quantities of food-based fortified lipid spreads (eg, lipid-based nutrient supplement). Such direct addition of micronutrients to foods is different from foods fortified in the preparatory processes, has the advantage that it does not require changing dietary practices, and has little effect on the taste of food. However, addition of micronutrient powders to prepared foods has characteristics akin to supplementation as opposed to foods fortified at source. Biofortification of food crops (fortification of food at source) is an alternative to more common fortification interventions and is rapidly advancing in technology with much success, particularly with regard to increasing iron, provitamin A, zinc, and folate contents in staple foods.¹¹²

Despite many limitations to establishing causality during assessment of food fortification programmes, several studies have reported outcomes. Fortification for children shows

significant benefits on serum micronutrient concentrations, which could indirectly be used to work out the population-level effect. A meta-analysis of multiple micronutrient fortification in children shows an increase in haemoglobin concentrations by 0.87 g/dL (95% CI 0.57–1.16) and reduced risk of anaemia by 57% (relative risk [RR] 0.43; 95% CI 0.26–0.71). The mean ferritin increase with fortification was 11.3 µg/L (95% CI 3.3–19.2) compared with control groups. Fortification also increased vitamin A serum concentrations compared with control groups (four studies, mean retinol increase 3.7 µg/dL, 95% CI 1.3–6.1).¹¹³ A meta-analysis of 60 trials showed that iron fortification of foods resulted in 41% reduction in the risks of anaemia (RR 0.59, 95% CI 0.48–0.71, $p < 0.001$) and a 52% reduction in iron deficiency (0.48, 0.38–0.62, $p < 0.001$).¹¹⁴ Other studies have also shown that use of vitamin D fortified bread increased serum 25-hydroxyvitamin D concentration as effectively as the cholecalciferol supplement in women.¹¹⁵ Zinc fortification has also shown significantly higher zinc concentrations in serum and erythrocytes and lower serum copper concentrations compared with a placebo group in preterm infants.¹¹⁶

Fortification has the greatest potential to improve the nutritional status of a population when implemented within a comprehensive nutrition strategy. Key issues to ensure a sustainable programme include: identification of the right food (accounting for bioavailability, interaction with food, availability, acceptability, and cost) and target population, ensuring quality of product, and consumption of sufficient quantity of the fortified foods.¹¹⁷ To accomplish these aims, there needs to be demand that is sustained through behaviour change communication at the consumer level and ready access to a sufficient supply of products that maintain standards set through legislative process from production to point-of-consumption. Government monitoring of compliance to standards and public-private partnerships are essential to ensure a competitive market for fortified products.¹¹⁸ Fortification seems to be a potentially effective strategy but evidence of benefits on morbidity and functional outcomes from large-scale programmes in developing countries is scarce.

population health. We reviewed relevant studies reporting the effect of financial incentives on coverage of health and nutrition interventions and behaviours targeting children younger than 5 years.¹²⁰ The affect of financial incentive programmes on five categories of interventions (breast-feeding practices, immunisation coverage, diarrhoea management, healthcare use, and other preventive strategies) was assessed. The review concluded that financial incentives have the potential to promote increased coverage of several important child health interventions, but the quality of evidence available was low. The more pronounced effects seemed to be achieved by programmes that directly removed user fees for access to health services.¹²⁰ Some indication of effect was also noted for programmes that conditioned financial incentives on participation in health education and attendance to health-care visits. Further

information on the benefit of such programmes for health and nutrition outcomes is provided in the accompanying report by Stuart Gillespie and colleagues.¹¹⁹

Community-based platforms for nutrition education and promotion

Community-based interventions to improve maternal, newborn, and child health are now widely recognised as important strategies to deliver key maternal and child survival interventions¹²¹ and have been shown to reduce inequities in childhood pneumonia and diarrhoea deaths.¹²² These interventions are delivered by health-care personnel or lay individuals, and implemented locally in homes, villages, or any defined community group. A full spectrum of promotive, preventive, and curative interventions can be delivered via community platforms,

including provision of basic antenatal, natal, and postnatal care; preventive essential newborn care; breastfeeding counselling; management and referral of sick neonates; development of skills in behaviour change communication; and community mobilisation strategies to promote birth and newborn care preparedness. For example, a review¹²³ of community-based packages of care suggested that these interventions can improve rates of facility births by 28% (RR 1.28, 95% CI 1.04–1.59) and result in a doubling of the rate of initiation of breastfeeding within 1 h (RR 2.25, 95% CI 1.70–2.97). Lewin and colleagues¹²⁴ reviewed 82 studies with lay health workers and showed moderate quality evidence of effect on initiation of breastfeeding (RR 1.36, 95% CI 1.14–1.61), any breastfeeding (1.24, 1.10–1.39), and exclusive breastfeeding (2.78, 1.74–4.44) when compared with usual care. Although much of the evidence from large-scale programmes using community health workers is of poor quality, process indicators and assessments do suggest that community health workers are able to implement many of these projects at scale, and have substantial potential to improve the uptake of child health and nutrition outcomes among difficult to reach populations.¹²⁵ It is important to underscore the

crucial importance of community engagement and buy-in to ensure effective community outreach programmes, behaviour change, and access.

Integrated management of childhood illnesses

WHO, in collaboration with UNICEF and other agencies, developed the Integrated Management of Childhood Illness (IMCI) strategy in the 1990s.¹²⁶ IMCI includes both curative and preventive interventions targeted at improvement of health practices at health facilities and at home. The strategy includes three components: improvements in case management; improvements in health systems; and improvements in family and community practices. Assessments of IMCI in Uganda, Tanzania, Bangladesh, Brazil, Peru, South Africa, China, Armenia, Nigeria, and Morocco have shown various benefits in health service quality, mortality reduction, and health-care cost savings.¹²⁷ In Tanzania, implementation of IMCI was associated with significant improvements in equity differentials for six child health indicators, with the largest improvements noted for stunting in children between 24 and 59 months of age.¹²⁸ Much the same findings were reported from Bangladesh, where implementation of IMCI was associated with a significant

Panel 4: Nutrition in emergencies

Irrespective of the underlying cause, humanitarian emergencies are often characterised by high and rising rates of severe acute malnutrition (SAM), moderate acute malnutrition (MAM), and micronutrient deficiencies in children (and sometimes adults). The foremost intent of nutrition-specific interventions in such situations is to prevent mortality, and involves management of wasting and resolution of specific nutrient deficiencies, and ensuring adequate food consumption. The humanitarian community largely agrees that emergency nutrition interventions have improved in the past 10–15 years in terms of coverage, scale of operations, reporting standards, and effectiveness (assessed by Sphere¹³³ and other standards of practice). Until the early 2000s, nutrition programming in emergencies was dominated by facility-based therapeutic care, targeted or blanket supplementary feeding, and provision of micronutrient supplements.^{134–136} More recently, the focus has widened, with attention being given to both short-term and longer-term concerns, and to a choice of actions from a more comprehensive range of interventions.¹³⁷ The options for effective management for both SAM and MAM in emergencies have improved in the past 10–15 years as products used have been improved and coverage has increased through community-based treatment. Potential alternatives to the use of food to address seasonal or emergency-driven peaks in wasting are being explored, including combinations of food plus cash, cash alone, or vouchers; however, cost-effectiveness studies of various strategies are scarce. There is evidence that in contexts in which markets have not been seriously disrupted, appropriate foods are readily accessible, and rates of

undernutrition are not dangerously high, alternatives or complements to food-based rations might be viable and potentially cost-effective.^{138,139}

Concern has also grown about the potential trade-offs between long-term versus short-term objectives of emergency nutrition interventions. Although life-saving actions are justifiably prioritised over the prevention of chronic diseases, food assistance programmes suitable for acute emergencies might be less appropriate for protracted situations.¹⁴⁰ This has important implications when thinking through seasonal blanket distribution of ready-to-use foods to prevent a worsening of levels of acute malnutrition. As a result of the difficulty of generating experimental data specific to programming in emergencies,¹⁴¹ the discipline has evolved relying less on randomised controlled trials and more on the sharing of lessons learned, which are used to inform technical or operational guidelines disseminated by WHO and UN bodies.^{142–144} Although practice must still be improved in many areas, and outcomes better documented, it remains crucially important to secure appropriate resources to support nutrition actions in this most challenging of disciplines and to assess outcomes for future learning. The nutritional status of individuals assessed and treated in emergency contexts overlaps substantially with non-emergency settings. Although high-quality programmatic research can and must help improve the design and outcome of effective emergency nutrition interventions, these interventions should be seen as entry points that support, rather than supplant, longer-term actions seeking to address underlying causes of poor nutrition.

increase in exclusive breastfeeding and comparatively faster reduction in the prevalence of stunting in children aged 24–59 months.¹²⁹

School-based delivery platforms

Many countries have school feeding programmes targeting children who are older than 5 years. The main purpose of such programmes is to provide incentives for school enrolment and evidence of nutrition benefits is scarce. A Cochrane review¹³⁰ of 18 relevant studies of the effectiveness of school feeding programmes in improving physical and psychosocial health for disadvantaged school pupils reported an increase in school attendance by 4–6 days annually and weight gains averaging 0.39 kg (95% CI 0.11–0.67) over 11 months and 0.71 kg (0.48–0.95) over 19 months. The results were inconclusive for height gain, so there must be caution that these programmes do not lead to obesity. A detailed discussion of school feeding programmes is provided in the accompanying report by Stuart Gillespie and colleagues.¹¹⁹ Notwithstanding the scarce evidence, schools offer an enormous opportunity for promotion of health and nutrition for older children and adolescents and could have an important role in future.

Child health days

Child health days have been introduced in weak health systems to rapidly enhance coverage of essential child

survival interventions. There are few robust assessments or reported experiences with child health days, which commonly include delivery of vitamin A supplements, immunisations, insecticide-treated nets, and deworming drugs. Available evidence suggests that these days can achieve greater coverage than stand-alone campaigns in previously low-coverage countries.¹³¹ A descriptive review¹³² of scale-up of child health days from 1999 to 2009 suggests that these days were more effective than stand-alone campaigns, provided that the number of interventions did not exceed four. The overall equity effect of these approaches are uncertain and further studies are needed to establish how best to integrate this approach within routine health-care services.

Delivery of nutrition interventions in humanitarian emergency settings

Delivery strategies for nutrition interventions in humanitarian emergencies necessitate a different approach to what might be deemed optimum in stable circumstances. In view of variability in the characteristics of emergencies and protracted population displacement, humanitarian emergencies might closely mirror situations of endemic malnutrition in food insecure settings. Hence prevention and health promotion strategies, such as breastfeeding and complementary feeding education and support, should also become essential parts of the packages of interventions in emergency contexts (panel 4^{133–144}).

Panel 5: Evidence for emerging interventions

Household air pollution

Household air pollution (HAP) from solid fuels used in simple stoves for cooking and heating, is recognised as a risk factor for several health outcomes with important consequences for child survival, including pneumonia,¹⁴⁶ low birthweight, and stillbirths.¹⁴⁷ A review of observational studies¹⁴⁸ shows significant risk reduction estimates for HAP for low birthweight (29%), stillbirth (34%), stunting (21%), and all-cause mortality (27%). Reduction of exposure to HAP could substantially reduce risk of several important outcomes for child survival. One randomised controlled trial in rural Guatemala,¹⁴⁹ with an improved stove intervention, reduced average exposure to indoor air pollution by 50% and resulted in a reduction in physician-diagnosed pneumonia (relative risk [RR] 0.84, 95% CI 0.63–1.13) although this difference was not statistically significant. However, this finding was supported by the results of an exposure-response analysis which showed a statistically significant reduction in the same outcome (0.82, 0.70–0.98). This intervention also resulted in a reduction in low birthweight (0.74, 95% CI 0.33–1.66), with babies weighing 89 g more (95% CI –27 to 204) than those in the control group.¹⁵⁰ A range of interventions, including both clean fuels and improved solid fuel stoves are available, but substantial challenges remain in achieving sustained use of low-cost low-emission technologies at scale in low-income households.

Maternal vitamin D supplementation

Vitamin D is an essential requirement of the body at any age. Vitamin D can be acquired through three main channels: through the skin via exposure to sunlight, from the diet, and from supplements or fortified foods. However, natural low-cost sources of dietary sources of vitamin D are very scarce. A systematic review¹⁵¹ assessing the association of vitamin D status in pregnancy, suggests that women with circulating 25-hydroxyvitamin D (25[OH]D) concentrations of less than 50 nmol/L in pregnancy have an increased risk of preeclampsia (odds ratio [OR] 2.09, 95% CI 1.50–2.90), gestational diabetes mellitus (1.38, 1.12–1.70), preterm birth (1.58, 1.08–2.31) and small-for-gestation age ([SGA] 1.52, 1.08–2.15). A long-term cohort study¹⁵² did not find any association of low maternal vitamin D concentrations with bone mineral content in late childhood. Similarly, a Cochrane review¹⁵³ assessed the effectiveness of vitamin D supplementation in pregnancy and revealed little evidence of benefits on functional pregnancy outcomes, although significant increase in serum vitamin D concentrations at term were noted and borderline reduction in low birthweight was reported in three trials (RR 0.48, 95% CI 0.23–1.01). The number of high-quality trials with maternal vitamin D supplementation is too small to draw conclusions on its usefulness and safety.

(Continues on next page)

(Continued from previous page)

Maternal zinc supplementation

A Cochrane review¹⁵⁴ suggests that zinc supplementation in pregnancy results in a 14% reduction in preterm birth (RR 0.86, 95% CI 0.76–0.97). This decrease was not accompanied by a similar reduction in stillbirths, neonatal death, SGA, or low birthweight. No subgroup differences were identified in women with low versus normal zinc nutrition levels or in women who complied with their treatment versus those who did not. We conclude that there is presently insufficient evidence for a beneficial role of isolated zinc supplementation in pregnancy.

Omega-3 fatty acid supplementation

Several reviews^{155–162} have been done to assess the effectiveness of maternal supplementation with omega-3 fatty acids during pregnancy and its effects on various outcomes including nutritional, morbidity, mortality, cognitive, and neurodevelopmental measures. Findings from these reviews, consisting of studies done in developed countries and of variable quality, suggest that marine omega-3 fatty acids administered in pregnancy reduce the rate of preterm birth and increase birthweight. However, a Cochrane review¹⁵⁵ suggests that there is not enough evidence to support the routine use of marine oil supplements or other prostaglandin precursors during pregnancy to reduce the risk of pre-eclampsia, preterm birth, low birthweight, or SGA. A review¹⁶³ of the intake of omega-3 and omega-6 fatty acids in low-income countries showed that the total omega-3 fatty acid supply was below the recommended intake range for infants and young children, and below the minimum recommended level for pregnant and lactating women, in the nine countries with the lowest gross domestic product. The review noted that supply of omega-3 fatty acids could be increased by using vegetable oils with higher alpha-linolenic acid and by increasing fish production through fish farming. Another review¹⁶⁴ on the effect of fatty acid status on immune function of children in low-income countries suggested that fatty acid interventions could yield immune benefits in children in poor settings, especially in non-breastfed children and in relation to inflammatory disorders, such as persistent enteropathy, although more trials are needed for a conclusive association.

Antenatal psychosocial assessment and mental health support

Stable maternal mental health during pregnancy is crucial for the development of the early mother–child relationship and for health. Although there is ample evidence of the link between maternal mental health and child health and growth,¹⁶⁵ there is insufficient evidence to support routine psycho-social screening for all pregnant women.¹⁶⁶ There is promising evidence that cognitive-behaviour therapy-based interventions provided by community health workers to pregnant women, can effectively reduce depression at 3 months post-partum (adjusted OR 0.22, 95% CI 0.14–0.36) and at 1-year follow-up (0.23, 0.15–0.36).¹⁶⁷ However, there was no effect on weight gain or linear growth in infancy. There is a need for further robust trials of maternal mental health interventions with longer term follow up.

Role of massage for promoting growth in preterm Infants

Preterm infants have been noted to benefit from massage therapy and the suggested mechanisms include increased vagal activity and gastric motility, which leads to increased concentrations of insulin and Insulin-like growth factor 1.¹⁶⁸ A Cochrane review¹⁶⁹ of the effect of massage in preterm infants showed that massage increased daily weight gain by 5 g, reduced the length of hospital stay by 4–5 days, and had a slight effect on development and weight gain at 4–6 months, although the evidence was of weak quality. A more recent review¹⁷⁰ of the effects of massage therapy for preterm infants showed that 5–10 days of moderate pressure massage, typically 15 min three-times daily, resulted in improved weight gain (mean for studies 28–48%) and bone density, and reduced length of hospital stay. Related evidence from studies of emollient therapy in preterm infants from the developing world suggest potential synergistic benefits of skin barrier protection, thermoregulation, and light massage.

Vitamin D supplementation in children

In view of the widespread deficiency of vitamin D and associated health consequences and rickets, preventive vitamin D supplementation to high-risk populations, including infants and toddlers, might be a useful strategy. A Cochrane review of vitamin D supplementation in children in at-risk populations is underway, and an existing review¹⁷¹ of postnatal supplementation shows relatively few studies assessing effects on bone density, growth, and other functional outcomes.

Zinc supplementation for treatment of newborn infections and childhood pneumonia

A Cochrane review¹⁷² suggests that zinc supplementation in addition to antibiotics in children with severe and non-severe pneumonia did not have a significant effect on clinical recovery or duration of hospital stay. Other recent studies show mixed effects across a range of severity of disease,^{173–176} showing the need for larger well-powered studies for the treatment of severe pneumonia with zinc in populations at-risk of deficiency. Two trials^{177,178} of adjunctive zinc supplementation in presumed serious infections in neonates and young infants show disparate findings, underscoring the need for further well-designed and adequately powered studies of zinc as an adjunct to the treatment of serious infections in infancy.

Lipid-based nutrient supplementation

Lipid-based nutrient supplements (LNS, in the form of vegetable oil, peanut butter, milk powder, sugar, vitamins, and minerals) are used in small quantities (20 g) to meet micronutrient requirements in children, in combination with a normal diet. Randomised controlled trials in Malawi^{179,180} and Ghana^{181,182} have shown significant benefits on iron status and linear growth. Further evidence of benefits and absence of adverse effects are needed to assess the feasibility of use of LNS in programme settings and randomised controlled trials are underway—three in Africa and one in Asia—which should provide more information.

Panel 6: Overview of the Lives Saved Tool (LiST)

To model the effect of scaling up the ten proven nutrition-specific interventions on the health of children we used the Lives Saved Tool (LiST). This model has been developed under the auspices of the Child Health Epidemiology Reference Group (CHERG) to allow users to estimate the effect of scaling up interventions on maternal and child health. The present version of the model is based on previous modelling exercises, including *The Lancet's* 2008 Maternal and Child Undernutrition series.^{4,122,145,184} A more detailed description of the LiST model is provided in appendix pp 3–7 and at the LiST website.

The LiST has been characterised as a linear, mathematical model that is deterministic.¹⁸⁵ It describes fixed associations between inputs and outputs that will produce the same outputs each time the model is run. In LiST the primary inputs are coverage of interventions and the outputs are changes in population level of risk factors (such as wasting or stunting rates, or birth outcomes such as prematurity or size at birth) and cause-specific mortality (neonatal, mortality in children aged 1–59 months, maternal mortality, and stillbirths). The association between an input (change in intervention coverage) with one or more outputs is specified in terms of the effectiveness of the intervention for reduction of the probability of that outcome. The outcome can be cause-specific mortality or a risk factor. The overarching assumption in LiST is that mortality rates and cause of death structure will not change except in response to changes in coverage of interventions. The model assumes that changes in distal variables, such as increase in income per person or mothers' education, will affect mortality by increasing coverage of interventions or reducing risk factors.

Figure 2 shows the linkages in LiST between risk factors, interventions, and mortality. For example, the input of multiple micronutrients consumed by pregnant women has an effect on birth outcomes. It directly affects the probability of a child being born small-for-gestational-age (SGA). A child born SGA will in turn have an increased risk of dying during the neonatal period and those who survive through the first month of life then have an increased risk of being stunted. Stunted children have higher risks of mortality from 1 to 59 months of age.

A second example is promotion of breastfeeding. Scaling up breastfeeding promotion will affect breastfeeding practice, shifting the distribution of mothers who exclusively, predominately, or partly breastfeed their child or do not breastfeed. Within the model, this intervention is examined separately for the first month of life and for the period of 1–5 months. Within each of the two time periods there is a different relative risk of dying of pneumonia and diarrhoea associated with each breastfeeding practice and an effect on diarrhoea incidence. So within the model, breastfeeding promotion has an effect on breastfeeding practices, which in turn has a direct effect on mortality in the neonatal and 1–23 month period. Additionally, there is an effect on diarrhoea incidence, which then has an effect on stunting rates and mortality.

The assumptions of the effects of the ten nutrition interventions we used in the modelling are shown in appendix pp 13–16. For each of the interventions we have also shown the 95% CIs around the estimates, which were used in the sensitivity analyses. The source and methods used to develop these assumptions and others used in the model are described in a series of reports.^{186–188}




Figure 2: Linkages between risk factors, interventions, and mortality in LiST

LiST=Lives Saved Tool. MMN=multiple micronutrients. BEP=balanced energy protein. SGA=small-for-gestational-age. SAM=severe acute malnutrition. MAM=moderate AM.

for inclusion in regular programmes. These interventions, which have possible effects on nutritional outcomes in women and children, include strategies to reduce household air pollution, maternal vitamin D supplementation, maternal zinc supplementation, omega 3 fatty acids supplementation in pregnancy, antenatal psychosocial assessment and cognitive behaviour therapy for depression, emollient and massage therapy for preterm infants, vitamin D supplementation in children, zinc therapy for pneumonia, and lipid-based nutrient supplements. Some of the existing evidence around these interventions is summarised in panel 5.^{146–182}

Modelling the effect of scaling up coverage of nutrition interventions in countries with the highest burden

We used the Lives Saved Tool (LiST) to model the potential effect on child health and mortality in 2012 of scaling up a set of ten nutrition-specific interventions that could affect stunting and severe wasting¹⁸³ (panel 6,^{4,122,145,184–188} figure 2). Although included in costing, we did not model the promotion and use of iodised salt. For modelling, we selected 34 countries with more than 90% of the burden

Emerging interventions that need further evidence

We also reviewed interventions that are not currently recommended but that have potential and future prospects




Figure 5: Effect of scale up of interventions on deaths in children younger than 5 years

Error bars are ranges. Promotion and use of iodised salt not modelled for mortality effect. SAM=severe acute malnutrition. MAM=moderate AM.

	Number of lives saved*	Cost per life-year saved†
Optimum maternal nutrition during pregnancy		
Maternal multiple micronutrient supplements to all	102 000	\$571 (398–1191)
Calcium supplementation to mothers at risk of low intake‡	(49 000–146 000)	
Maternal balanced energy protein supplements as needed		
Universal salt iodisation‡		
Infant and young child feeding		
Promotion of early and exclusive breastfeeding for 6 months and continued breastfeeding for up to 24 months	221 000	\$175 (132–286)
Appropriate complementary feeding education in food secure populations and additional complementary food supplements in food insecure populations	(135 000–293 000)	
Micronutrient supplementation in children at risk		
Vitamin A supplementation between 6 and 59 months age	145 000	\$159 (106–766)
Preventive zinc supplements between 12 and 59 months of age	(30 000–216 000)	
Management of acute malnutrition		
Management of moderate acute malnutrition	435 000	\$125 (119–152)§
Management of severe acute malnutrition	(285 000–482 000)	

Data are number (95% CI) or cost in 2010 international dollars (95% CI). *Effect of each of package when all four packages are scaled up at once. †Cost per life-year saved assumes that a life saved of a child younger than 5 years saves on average 59 life-years, based on WHO data (2011¹⁸⁹) that life expectancy at birth on average in low-income countries is 60, and that most deaths of children younger than 5 years occur in the first year of life. To convert to cost per discounted life-year saved multiply these estimates by 59/32 (ie, 1.84). ‡Intervention has effect on maternal or child morbidity, but no direct effect on lives saved. §Cost per life-year saved by management of severe acute malnutrition only, costs for supplementary feeding for moderate acute malnutrition are currently unavailable.

Table 5: Effect of packages of nutrition interventions at 90% coverage

Implementation of nutrition-specific packages of care

We also assessed the potential effect of nutrition-specific packages of care by scaling up these interventions to 90% coverage. Four packages were assessed for effect on child survival: optimum maternal nutrition during pregnancy (maternal multiple micronutrients, use of iodised salt, calcium, and balanced energy protein supplementation), an infant and young child nutrition package (breastfeeding promotion and appropriate complementary feeding education or provision), micronutrient supplementation (preventive zinc and vitamin A supplementation), and management of acute malnutrition (management of MAM, management of SAM). Analysis

of these nutrition-specific packages showed that the most lives could be saved by the therapeutic feeding for severe acute malnutrition, followed by the infant and young child nutrition package (table 5¹⁸⁹).

Can these interventions promote equitable access?

To assess the potential benefit of community-based delivery strategies on reaching and engaging poor and marginalised populations, we assessed the effect of community-based promotion and delivery of seven nutrition-specific interventions (multiple micronutrient supplementation in pregnancy, promotion of breastfeeding, appropriate complementary feeding, management of SAM, vitamin A supplementation, preventive zinc supplementation, and treatment of diarrhoea with zinc) across various wealth quintiles in three target countries—Pakistan, Bangladesh, and Ethiopia (figure 6). Baseline data were stratified by wealth quintiles by reanalysing the most recently available Demographic Health Survey for each country. Since no recent estimates existed for cause-specific mortality across wealth quintiles for Bangladesh and Ethiopia, we used LiST to recompute the cause of death structure using the procedures described by Amouzou and colleagues.¹⁹⁰ For Pakistan, we used the recent national verbal autopsy study¹⁹¹ for the cause of death structure and distribution of deaths by asset quintiles from a recent analysis.¹⁹² As shown in figure 6 and appendix p 26, the effect of this scale up is greatest in the poorest quintiles, suggesting that scaling up these interventions through community-based approaches would not only reduce the overall burden of childhood mortality but also substantially reduce existing disparities in access and mortality.

Cost analysis

We used a so-called ingredients approach to work out the cost of nutrition interventions, based on the UN One Health Tool,¹⁹³ which allows for regional variation due to personnel costs. We constructed cost estimates as additions to existing antenatal, postnatal, and standard infant




Figure 6: Equity analysis showing effect of scale up of nutrition interventions on proportion of deaths of children younger than 5 years averted in Pakistan, Bangladesh, and Ethiopia
Error bars are ranges.

visits as part of WHO's Expanded Program on Immunisation, plus five stand-alone nutrition visits between 6 and 35 months of age. The few interventions targeted at children between 36 and 59 months of age were assumed to be delivered opportunistically (at clinic visits, or during outreach visits for younger siblings). The base delivery platform assumed was outreach programmes for sub-Saharan Africa, and primary health-care clinics elsewhere (appendix pp 27–28 provides details). We compared the unit costs from the ingredients method with actual costs as used in the Scaling up Nutrition (SUN) costing.¹⁹⁴ Although the ingredients method allows greater detail than an actual costs method such as SUN for planning purposes, the comparison to actual costs serves as a useful check on the appropriateness of assumptions made. Costs were estimated for ten nutrition interventions (one population-wide, three in

	Cost
Salt iodisation	\$68
Multiple micronutrient supplementation in pregnancy (includes iron-folate)	\$472
Calcium supplementation in pregnancy	\$1914
Energy-protein supplementation in pregnancy	\$972
Vitamin A supplementation in childhood	\$106
Zinc supplementation in childhood	\$1182
Breastfeeding promotion	\$653
Complementary feeding education	\$269
Complementary food supplementation	\$1359
SAM management	\$2563
Total	\$9559

Data are 2010 international dollars, millions.

Table 6: Total additional annual cost of achieving 90% coverage with nutrition interventions, excluding management of moderate acute malnutrition, in 34 countries with more than 90% of the burden

pregnancy, and six after birth; intervention definitions and assumptions are provide in appendix pp 29–30). Costs were estimates for a set of ten nutrition interventions (including use of iodised salt). Unit costs for management of MAM could not be obtained.

We calculated unit costs separately for WHO sub-regions (appendix p 31).¹⁹⁵ Unit costs were higher in Africa compared with elsewhere, because of higher labour costs and the extra travel time required for delivery using outreach (associated with lower population density in many areas, and also the lower coverage of primary care facilities). The unit costs for interventions were much the same between the ingredients and SUN approaches, allowing for some difference in interventions based on updated recommendations.

Our analysis shows that the estimated total additional cost involved to achieve 90% coverage of the population in need in the 34 focus countries with the selected set of ten nutrition interventions is Int\$9.6 billion per annum (table 6). Of this \$9.6 billion, \$3.7 billion (39%) is for micronutrient interventions, \$0.9 billion (9%) for educational interventions, and \$2.6 billion (27%) for SAM management. The amount required for provision of supplementary food for pregnant women and for children aged 6–23 months in poor households (those with <\$1.25 per person per day) constitutes the remaining \$2.3 billion (24%). When these costs are broken down by region, \$3.4 billion is needed in the 20 countries included from sub-Saharan Africa, \$4.8 billion in the four in south Asia plus Myanmar (Burma), \$1.0 billion for the six in eastern Mediterranean, and \$0.5 billion for the three remaining countries (Vietnam and the Philippines in western Pacific region, plus Guatemala; appendix p 32). The \$9.6 billion estimate for the nutrition interventions is lower than the 2008 SUN estimate of \$11.8 billion.¹⁹⁴ The SUN figure included \$1.2 billion for capacity-building and monitoring and assessment, which we

excluded from the present analysis because we do not have a mechanism to allocate this cost by region or country and category.

There are differences in the details of our results compared with the earlier SUN estimates. Unit costs are similar but not identical. The list of focus countries is likewise similar but not identical (using 2005 data, Turkey, Peru, Cambodia, and Burundi were included in the list of countries with 90% of the world's stunted children, but not with 2010 data; Rwanda and Chad entered this list with 2010 data). Some interventions are excluded from the new total (deworming, therapeutic zinc for diarrhoea), whereas some new ones are included (calcium supplements and balanced energy protein supplements in pregnancy). Complementary food supplementation is targeted only to the 6–23 month age group in this new analysis. Population in need has changed since the SUN estimates were calculated, with changes in coverage of some interventions (notably, management of SAM has begun to scale up).

Discussion

This update of nutrition interventions differs from past exercises in several ways. First, we included a wider range of nutrition-specific interventions and applied more stringent assessment criteria, using the Grades of Recommendation Assessment, Development and Evaluation system and Child Health Epidemiology Reference Group criteria for most interventions.⁵ Second, in view of emerging evidence of the importance of maternal nutrition, SGA, and early stunting,¹ we specifically focused on interventions that might affect prevalence and outcomes in SGA births and early stunting. We also reviewed and modelled a range of delivery strategies and platforms and specifically explored the potential of reaching poor and disadvantaged populations through community platforms and outreach services, an approach used to assess the effect of interventions to address childhood diarrhoea and pneumonia.¹²² Finally, the LiST model was substantially updated to include age-specific effects and the inter-relationship of new interventions and their effect on maternal and child undernutrition, a much more complex exercise than what was undertaken previously.⁴ To undertake this exercise we substantively updated LiST in a way that more accurately captures the role of undernutrition and the effect of proven interventions on maternal and child health.

Several limitations should be recognised in considering our findings. A large proportion of the evidence on interventions is still derived from efficacy trials as opposed to effectiveness studies and hence variations exist in estimates of effect size for various interventions. Few robust assessments have been done in programme settings and available data from observational studies do not permit ready assessment of intervention effectiveness. There are also very few studies that report morbidity and neurodevelopmental outcomes. We reviewed the available evidence of effects on neurodevelopmental outcomes

from studies of maternal and child nutrition interventions and identified little data with evaluation methodologies for assessment. We are therefore greatly limited in the inferences that can be drawn on neurodevelopment and long-term outcomes from nutrition interventions.

Notwithstanding these limitations, our estimates of the effect of nutrition-specific interventions, though more conservative than previous findings, still suggest great benefits from a core set of interventions delivered antenatally or postnatally. Our assessments of benefits from interventions to reduce SGA are hindered by the limited range of interventions in pregnancy. Even though antenatal care services offer a unique opportunity for maternal screening and interventions, the difficulty of reaching women early enough in pregnancy is a major limitation in ensuring adequate uptake of interventions for a reasonable length of time. In parts of the world with high rates of maternal malnutrition, micronutrient deficiencies, and SGA births, these factors remain major determinants of stunting in early childhood. This finding underscores the need to address determinants of undernutrition early in the lifecycle through appropriate strategies, such as enhancing adolescent nutrition and family planning to delay the age of first pregnancy or increase spacing between births.¹⁹⁶ Achieving high coverage of multiple micronutrient supplementation in pregnancy offers a new avenue to reduce SGA births and their consequences for mortality and growth in early childhood. The absence of appreciation of the crucial links of maternal and fetal nutrition to fertility and repeated pregnancies has been a major barrier in targeting of interventions to address these factors appropriately.

Although the overall effect on stunting alleviation seems modest, the rate of decline suggested from the package of nutrition-specific interventions is plausible and within the broad range of observed effects across countries. A review of global stunting trends by Stevens and colleagues¹⁹⁷ showed average rates of reduction in stunting in the best performing countries ranging from 21–42% over the past decade, broadly consistent with what our model predicts from scaling up a core set of nutrition-specific interventions. Importantly, the countries that have made tremendous strides in improving nutrition and health outcomes (such as Brazil, China, Saudi Arabia, Kuwait, and Chile) have implemented nutrition-specific interventions, but also have been settings with exceptional economic growth, and investments in nutrition-sensitive interventions to address population health, education, and social sector development. Much the same conclusions were drawn by UNICEF in a report on undernutrition.¹⁹⁸

A major advance on our previous review of interventions is the addition of delivery platforms that allow us to assess strategies to reach populations who are not being reached currently. Our findings suggest that community platforms offer a unique opportunity to engage and reach poor and difficult to access populations through communication and outreach strategies. These

strategies could also lead to potential integration of nutrition with maternal, newborn, and child health interventions. Since several countries are investing in community health worker programmes to address maternal, newborn, and child health,¹²⁵ much potential exists for scaling up nutrition promotion and therapeutic interventions through such platforms and hence integrating the two at point-of-service delivery. This integration could also help achieve reductions in inequities in the short term as has been noted by universal scaling up of selected maternal and child survival interventions.¹⁹⁹ However, importantly, implementation of such programmes involves unique combinations and sequencing of health system policies, actions, and advocacy. Community-based nutrition programmes need meticulous planning, a rights-based framework for engagement of communities and other sectors, and piloting. Other health system pillars are crucial to success, including training and support for community health workers, strengthening of the supply chain, simplified information systems, monitoring, and regular feedback.

The model used in this review estimates feasible reductions in mortality and stunting with enhanced investments. The same investments that can achieve these results will also lead to other improvements in cognitive and socioemotional development. Although these outcomes are not included in the model, partly because the costing database differs (ie, LiST mainly addresses mortality), substantial evidence from a range of models and longitudinal studies confirms that the benefits in terms of overall development on human capacity are appreciable.

In terms of cost, an annual outlay of an additional \$9.6 billion to bring to scale a range of nutrition interventions that would save nearly 1 million lives is reasonable since many interventions would be scaled up from negligible coverage rates. The cost per (discounted) life-year saved is about \$370 for a set of interventions that could effectively deliver optimum nutrition to pregnant women, infants, and children, and manage SAM. These figures suggest that the nutrition interventions are well within the cost-effectiveness benchmark (less than three-times per person income) for all countries. More than half the \$9.6 billion is accounted for by two large countries that could rely heavily on domestic resources (India and Indonesia). Consumables (whether drugs, or other items such as for transport or administration) account for slightly less than half the \$9.6 billion, and all but the poorest countries can be expected to cover most of the expenditures on personnel; \$3–4 billion from external donors could make a substantial difference to child nutrition. What proportion of development assistance for health is earmarked for nutrition is unclear. Global tracking data from the Institute for Health Metrics and Evaluation were unable to disaggregate nutrition-related funding from the annual funding of \$5.17 billion for

maternal, newborn, and child health programmes.²⁰⁰ The Countdown collaboration estimates nutrition-specific funding for the same year (2012) at \$324.5 million.²⁰¹ Much funding for nutrition-relevant programmes probably overlaps with existing programmes for maternal, newborn, and child health and health systems strengthening and there might also be potential synergies, making it possible to share costs.

The evidence from carefully conducted cohort studies²⁰² of benefits of higher birthweight and early linear growth on education and improved health outcomes is convincing and consistent with the overall message from our review and modelling exercise. As the world moves towards the post-2015 development agenda, it is important to draw attention to the unfinished agenda of maternal and child undernutrition and to the emerging issues of obesity. Our review reconfirms the existence of feasible and low cost evidence-based interventions and the fact that coverage rates for many of these interventions remain poor²⁰³ and for some, non-existent. In view of the importance of fetal nutrition and poverty alleviation strategies to reach those in greatest need, priority must be given to scaling up nutrition specific and sensitive interventions in some of the highest burden countries. At the same time, in view of the increasing importance of non-communicable diseases, concerted efforts must be made to develop and implement interventions to reduce the risk of obesity.

Contributors

ZAB conceptualised the review in consultation with the coordinators (PW, AL, SH, and REB) and wrote the first draft of the paper with substantial inputs from JKD, NW, YT, and AR developed the modification of LiST for assessment of effect and equity. Costing for selected interventions was done by MFG and SH. VW contributed to the scientific literature search, screening, collection, and analysis of data for the deworming review. LL led the review of severe and moderate acute malnutrition. KW led the obesity prevention review and contributed to the severe and moderate acute malnutrition reviews. CM and SZ oversaw the obesity reviews. BAH assessed the effect of maternal multiple micronutrient supplements, neonatal vitamin A supplementation, and deworming in pregnant women. ZL contributed to reviews of complementary feeding strategies and community platforms. RAS contributed to reviews of micronutrient powders and breastfeeding. AI contributed to the reviews of calcium, balanced protein supplementation in pregnancy, and vitamin A supplementation. All authors and members of the review groups (below) saw successive drafts of the paper and provided input. ZAB finalized the paper and is the overall guarantor.

The Lancet Nutrition Interventions Review Group

Zulfiqar A Bhutta, Arjumand Rizvi, Jai K Das, Rehana A Salam, Aisha Yousafzai (Aga Khan University, Pakistan), Zohra S Lassi (University of Adelaide, Australia), Lindsey Lenters, Ceilidh McPhail, Kerri Wazny, Michelle F Gaffey, Stanley Zlotkin (Hospital for Sick Children, Canada), Aamer Imdad (SUNY Upstate Medical University, USA), Batool Azra Haider (Harvard School of Public Health, USA), Vivian Welch (University of Ottawa, Canada), Reynaldo Martorell (Emory University, USA), Robert E Black, Neff Walker, Yvonne Tam (Johns Hopkins University, USA), Tahmeed Ahmad (International Center for Diarrheal Diseases Research, Bangladesh).

Maternal and Child Nutrition Study Group

Robert E Black (Johns Hopkins Bloomberg School of Public Health, USA), Cesar Victora (Universidade Federal de Pelotas, Brazil), Susan Walker (The University of the West Indies, Jamaica),

Harold Alderman (International Food Policy Research Institute, USA), Zulfiqar A Bhutta (Aga Khan University, Pakistan), Stuart Gillespie (International Food Policy Research Institute, USA), Lawrence Haddad (Institute of Development Studies, UK), Susan Horton (University of Waterloo, Canada), Anna Larrey (University of Ghana, Ghana), Venkatesh Mannar (The Micronutrient Initiative, Canada), Marie Ruel (International Food Policy Research Institute, USA), Patrick Webb (Tufts University, USA)

Series Advisory Committee

Marc Van Ameringen (Gain Health Organization, Switzerland), Mandana Arabi (New York Academy of Sciences, USA), Shawn Baker (Helen Keller International, USA), Martin Bloem (United Nations World Food Programme, Italy), Francesco Branca (WHO, Switzerland), Leslie Elder (The World Bank, USA), Erin McLean (Canadian International Development Agency, Canada), Carlos Monteiro (University of São Paulo, Brazil), Robert Mwandime (Makerere School of Public Health, Uganda), Ellen Piwoz (Bill & Melinda Gates Foundation, USA), Werner Schultink (UNICEF, USA), Lucy Sullivan (1000 Days, USA), Anna Taylor (Department for International Development, UK), Derek Yach (The Vitality Group, USA). The Advisory Committee provided advice in a meeting with Series Coordinators for each paper at the beginning of the process to prepare the Series and in a meeting to review and critique the draft reports.

Conflicts of interest

REB serves on the Boards of the Micronutrient Initiative, Vitamin Angels, the Child Health and Nutrition Research Initiative, and the Nestlé Creating Shared Value Advisory Committee. VM serves on the Nestlé Creating Shared Value Advisory Committee. The other authors declare that they have no conflicts of interest. As corresponding author Zulfiqar A Bhutta states that he had full access to all data and final responsibility to submit for publication.

Acknowledgments

Funding for the preparation of the Series was provided to the Johns Hopkins Bloomberg School of Public Health through a grant from the Bill & Melinda Gates Foundation. The sponsor had no role in the analysis and interpretation of the evidence or in writing the paper and the decision to submit for publication. We thank the Aga Khan University, Karachi, Pakistan and the Program for Global Pediatric Research, Global Child Health, Hospital for Sick Children (Toronto, ON, Canada) for support to ZAB during the course of this work.

References

- Black RE, Victora CG, Walker SP, and the Maternal and Child Nutrition Study Group. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* 2013; published online June 6. [http://dx.doi.org/10.1016/S0140-6736\(13\)60937-X](http://dx.doi.org/10.1016/S0140-6736(13)60937-X).
- Liu L, Johnson HL, Cousens S, et al, for the Child Health Epidemiology Reference Group of WHO and UNICEF. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet* 2012; **379**: 2151–61.
- Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**: 2224–60.
- Bhutta ZA, Ahmed T, Black RE, et al, for the Maternal and Child Undernutrition Group. What works? Interventions for maternal and child undernutrition and survival. *Lancet* 2008; **371**: 417–40.
- Walker N, Fischer-Walker C, Bryce J, Bahl R, Cousens S. Standards for CHERG reviews of intervention effects on child survival. *Int J Epidemiol* 2010; **39** (suppl 1): i21–31.
- WHO, The Partnership for Maternal, Newborn and Child Health. Reaching child brides. 2012. http://www.who.int/pmnch/topics/part_publications/knowledge_summary_22_reaching_child_brides/en/ (accessed March 25, 2013).
- Ronnenberg AG, Wood RJ, Wang X, et al. Preconception hemoglobin and ferritin concentrations are associated with pregnancy outcome in a prospective cohort of Chinese women. *J Nutr* 2004; **134**: 2586.
- Mehra S, Agrawal D. Adolescent health determinants for pregnancy and child health outcomes among the urban poor. *Indian Pediatr* 2004; **41**: 137.
- Haldre K, Rahu K, Karro H, Rahu M. Is a poor pregnancy outcome related to young maternal age? A study of teenagers in Estonia during the period of major socio-economic changes (from 1992 to 2002). *Eur J Obstet Gynecol Reprod Biol* 2007; **131**: 45–51.
- Paranjothy S, Broughton H, Adappa R, Fone D. Teenage pregnancy: who suffers? *Arch Dis Child* 2009; **94**: 239.
- WHO. Adolescent pregnancy: unmet needs and undone deeds. Geneva: World Health Organization, 2007.
- UNFPA. Marrying too young: end child marriage. New York, NY: UN Population Fund, 2012.
- Conde-Agudelo A, Rosas-Bermudez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. *JAMA* 2006; **295**: 1809.
- Conde-Agudelo A, Rosas-Bermudez A, Castaño F, Norton MH. Effects of birth spacing on maternal, perinatal, infant, and child health: a systematic review of causal mechanisms. *Stud Fam Plann* 2012; **43**: 93–114.
- WHO. Meeting report. Meeting to develop a global consensus on preconception care to reduce maternal and childhood mortality and morbidity; Geneva; Feb 6–7, 2012.
- Rosenthal AN, Paterson Brown S. Is there an incremental rise in the risk of obstetric intervention with increasing maternal age? *BJOG* 1998; **105**: 1064–69.
- Carolan M. The graying of the obstetric population: implications for the older mother. *J Obstet Gynecol Neonatal Nurs* 2003; **32**: 19–27.
- Alkema L, Kantorova V, Menozzi C, Biddlecom A. National, regional, and global rates and trends in contraceptive prevalence and unmet need for family planning between 1990 and 2015: a systematic and comprehensive analysis. *Lancet* 2013; published online March 13. [http://dx.doi.org/10.1016/S0140-6736\(12\)62204-1](http://dx.doi.org/10.1016/S0140-6736(12)62204-1).
- De-Regil LM, Fernandez-Gaxiola AC, Dowswell T, Pena-Rosas JP. Effects and safety of periconceptional folate supplementation for preventing birth defects. *Cochrane Database Syst Rev* 2010; **10**: CD007950.
- Lassi ZS, Salam RA, Haider BA, Bhutta ZA. Folic acid supplementation during pregnancy for maternal health and pregnancy outcomes. *Cochrane Database Syst Rev* 2013; **3**: CD006896.
- Fernandez-Gaxiola AC, De-Regil LM. Intermittent iron supplementation for reducing anaemia and its associated impairments in menstruating women. *Cochrane Database Syst Rev* 2011; **12**: CD009218.
- Pena-Rosas JP, De-Regil LM, Dowswell T, Viteri FE. Daily oral iron supplementation during pregnancy. *Cochrane Database Syst Rev* 2012; **12**: CD004736.
- Haider BA, Bhutta ZA. Multiple-micronutrient supplementation for women during pregnancy. *Cochrane Database Syst Rev* 2012; **11**: CD004905.
- Imdad A, Bhutta ZA. Effects of calcium supplementation during pregnancy on maternal, fetal and birth outcomes. *Paediatr Perinat Epidemiol* 2012; **26**: 138–52.
- Zimmermann MB. The effects of iodine deficiency in pregnancy and infancy. *Paediatr Perinat Epidemiol* 2012; **26** (suppl 1): 108–17.
- Imdad A, Bhutta ZA. Maternal nutrition and birth outcomes: effect of balanced protein-energy supplementation. *Paediatr Perinat Epidemiol* 2012; **26** (suppl 1): 178–90.
- Pena-Rosas JP, De-Regil LM, Dowswell T, Viteri FE. Intermittent oral iron supplementation during pregnancy. *Cochrane Database Syst Rev* 2012; **7**: CD009997.
- WHO. Guideline: daily iron and folic acid supplementation in pregnant women. Geneva: World Health Organization, 2012.
- Ronsmans C, Fisher DJ, Osmond C, Margetts BM, Fall CH. Multiple micronutrient supplementation during pregnancy in low-income countries: a meta-analysis of effects on stillbirths and on early and late neonatal mortality. *Food Nutr Bull* 2009; **30**: S547–55.
- Vaidya A, Saville N, Shrestha BP, Costello AM, Manandhar DS, Osrin D. Effects of antenatal multiple micronutrient supplementation on children's weight and size at 2 years of age in Nepal: follow-up of a double-blind randomised controlled trial. *Lancet* 2008; **371**: 492–99.

- 31 West KP Jr, Shamim AA, Labrique AB, et al. Efficacy of antenatal multiple micronutrient (MM) vs iron-folic acid supplementation in improving gestational and postnatal viability in rural Bangladesh: the JiVitA-3 Trial. *FASEB J* 2013; 27: 5225 (abstr).
- 32 Srinivas SK, Edlow AG, Neff PM, Sammel MD, Andrela CM, Elovitz MA. Rethinking IUGR in preeclampsia: dependent or independent of maternal hypertension? *J Perinatol* 2009; 29: 680–84.
- 33 Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol* 2009; 33: 130–37.
- 34 Hofmeyr GJ, Lawrie TA, Atallah AN, Duley L. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database Syst Rev* 2010; 8: CD001059.
- 35 Wu T, Liu GJ, Li P, Clar C. Iodised salt for preventing iodine deficiency disorders. *Cochrane Database Syst Rev* 2006; 3: CD003204.
- 36 Kulier R, de Onis M, Gulmezoglu AM, Villar J. Nutritional interventions for the prevention of maternal morbidity. *Int J Gynaecol Obstet* 1998; 63: 231–46.
- 37 de Onis M, Villar J, Gulmezoglu M. Nutritional interventions to prevent intrauterine growth retardation: evidence from randomized controlled trials. *Eur J Clin Nutr* 1998; 52 (suppl 1): S83–93.
- 38 Kramer MS, Kakuma R. Energy and protein intake in pregnancy. *Cochrane Database Syst Rev* 2003; 4: CD000032.
- 39 McDonald SJ, Middleton P. Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes. *Cochrane Database Syst Rev* 2009; 2: CD004074.
- 40 Rabe H, Diaz-Rossello JL, Duley L, Dowswell T. Effect of timing of umbilical cord clamping and other strategies to influence placental transfusion at preterm birth on maternal and infant outcomes. *Cochrane Database Syst Rev* 2011; 8: CD003248.
- 41 Puckett RM, Offringa M. Prophylactic vitamin K for vitamin K deficiency bleeding in neonates. *Cochrane Database Syst Rev* 2010; 4: CD002776.
- 42 Darlow BA, Graham PJ. Vitamin A supplementation to prevent mortality and short and long-term morbidity in very low birthweight infants. *Cochrane Database Syst Rev* 2007; 4: CD000501.
- 43 Haider BA, Bhutta ZA. Neonatal vitamin A supplementation for the prevention of mortality and morbidity in term neonates in developing countries. *Cochrane Database Syst Rev* 2011; 10: CD006980.
- 44 Moore ER, Anderson GC, Bergman N, Dowswell T. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev* 2012; 5: CD003519.
- 45 Conde-Agudelo A, Belizan JM, Diaz-Rossello J. Kangaroo mother care to reduce morbidity and mortality in low birthweight infants. *Cochrane Database Syst Rev* 2011; 3: CD002771.
- 46 WHO. Meeting report. Technical consultation on vitamin a in newborn health: mechanistic studies; Geneva; Dec 1–3, 2009.
- 47 Dyson L, McCormick FM, Renfrew MJ. Interventions for promoting the initiation of breastfeeding. *Cochrane Database Syst Rev* 2005; 2: CD001688.
- 48 Cai X WT, Brown DW. Global trends in exclusive breastfeeding. *Int Breastfeed J* 2012; 7: 12.
- 49 Edmond KM, Zandoh C, Quigley MA, Amenga-Etego S, Owusu-Agyei S, Kirkwood BR. Delayed breastfeeding initiation increases risk of neonatal mortality. *Pediatrics* 2006; 117: e380–6.
- 50 Mullany LC, Katz J, Li YM, et al. Breast-feeding patterns, time to initiation, and mortality risk among newborns in southern Nepal. *J Nutr* 2008; 138: 599–603.
- 51 Debes AK, Kohli A, Walker N, Edmond K, Mullany LC. Time to initiation of breastfeeding and neonatal mortality and morbidity: a systematic review. *BMC Public Health* (submitted).
- 52 Imdad A, Yakoob MY, Bhutta ZA. Effect on breastfeeding promotion interventions on breastfeeding rates, with special focus on developing countries. *BMC Public Health* 2011; 11 (suppl 3): S24.
- 53 Haroon S, Das JK, Salam RA, Bhutta ZA. Breastfeeding promotion interventions and breastfeeding practices: a systematic review. *BMC Public Health* (in press).
- 54 Lassi ZS, Zahid GS, Das JK, Bhutta ZA. Impact of complementary food and education on complementary food on growth and morbidity of children less than 2 years of age in developing countries: a systematic review. *BMC Public Health* (in press).
- 55 Imdad A, Herzer K, Mayo-Wilson E, Yakoob MY, Bhutta ZA. Vitamin A supplementation for preventing morbidity and mortality in children from 6 months to 5 years of age. *Cochrane Database Syst Rev* 2010; 12: CD008524.
- 56 De-Regil LM, Jefferds MED, Sylvetsky AC, Dowswell T. Intermittent iron supplementation for improving nutrition and development in children under 12 years of age. *Cochrane Database Syst Rev* 2011; 12: CD009085.
- 57 Sachdev H, Gera T, Nestel P. Effect of iron supplementation on mental and motor development in children: systematic review of randomised controlled trials. *Public Health Nutr* 2005; 8: 117–32.
- 58 Allen LH, Peerson JM, Olney DK. Provision of multiple rather than two or fewer micronutrients more effectively improves growth and other outcomes in micronutrient-deficient children and adults. *J Nutr* 2009; 139: 1022–30.
- 59 Salam RA, MacPhail C, Das JK, Bhutta ZA. Effectiveness of micronutrient powders (MNP) in women and children. *BMC Public Health* (in press).
- 60 Imdad A, Bhutta ZA. Effect of preventive zinc supplementation on linear growth in children under 5 years of age in developing countries: a meta-analysis of studies for input to the lives saved tool. *BMC Public Health*. 2011; 11 (suppl 3): S22.
- 61 Yakoob MY, Theodoratou E, Jabeen A, et al. Preventive zinc supplementation in developing countries: impact on mortality and morbidity due to diarrhea, pneumonia and malaria. *BMC Public Health* 2011; 11 (suppl 3): S23.
- 62 Gogia S, Sachdev HS. Zinc supplementation for mental and motor development in children. *Cochrane Database Syst Rev* 2012; 12: CD007991.
- 63 Abdulwadud OA, Snow ME. Interventions in the workplace to support breastfeeding for women in employment. *Cochrane Database Syst Rev* 2012; 3: CD006177.
- 64 WHO. Meeting report. Informal meeting to review and develop indicators for complementary feeding. Washington DC: World Health Organization, 2002.
- 65 Ruel MT, Menon P. Child feeding practices are associated with child nutritional status in Latin America: innovative uses of the demographic and health surveys. *J Nutr* 2002; 132: 1180–87.
- 66 Arimond M, Ruel MT. Dietary diversity is associated with child nutritional status: evidence from 11 demographic and health surveys. *J Nutr* 2004; 134: 2579–85.
- 67 Marriott BP, White A, Hadden L, Davies JC, Wallingford JC. World Health Organization (WHO) infant and young child feeding indicators: associations with growth measures in 14 low-income countries. *Matern Child Nutr* 2012; 8: 354–70.
- 68 Dewey KG, Adu-Afaruwah S. Systematic review of the efficacy and effectiveness of complementary feeding interventions in developing countries. *Matern Child Nutr* 2008; 4 (suppl 1): 24–85.
- 69 Awasthi S, Peto R, Read S, et al, and the DEVTA (Deworming and Enhanced Vitamin A) team. Vitamin A supplementation every 6 months with retinol in 1 million pre-school children in north India: DEVTA, a cluster-randomised trial. *Lancet* 2013; 381: 1469–77.
- 70 Sazawal S, Black RE, Ramsan M, et al. Effects of routine prophylactic supplementation with iron and folic acid on admission to hospital and mortality in preschool children in a high malaria transmission setting: community-based, randomised, placebo-controlled trial. *Lancet* 2006; 367: 133–43.
- 71 WHO. Guideline: intermittent iron supplementation in preschool and school-age children. Geneva: World Health Organization, 2011.
- 72 Neufeld LM, Ramakrishnan U. Multiple micronutrient interventions during early childhood: moving towards evidence-based policy and program planning. *J Nutr* 2011; 141: 2064–65.
- 73 Eilander A, Gera T, Sachdev HS, et al. Multiple micronutrient supplementation for improving cognitive performance in children: systematic review of randomized controlled trials. *Am J Clin Nutr* 2010; 91: 115–30.
- 74 Gera T, Sachdev HP. Effect of iron supplementation on incidence of infectious illness in children: systematic review. *BMJ* 2002; 325: 1142.
- 75 Soofi SB, Cousens S, Iqbal SP, et al. Effect of provision of daily zinc and iron with several micronutrients on growth and morbidity among young children in Pakistan: a cluster-randomised trial. *Lancet* 2013; published online April 18. [http://dx.doi.org/10.1016/S0140-6736\(13\)60437-7](http://dx.doi.org/10.1016/S0140-6736(13)60437-7).

- 76 Brown KH, Peerson JM, Baker SK, Hess SY. Preventive zinc supplementation among infants, preschoolers, and older prepubertal children. *Food Nutr Bull* 2009; **30** (suppl 1): S1–40.
- 77 Cairncross S, Hunt C, Boisson S, et al. Water, sanitation and hygiene for the prevention of diarrhoea. *Int J Epidemiol* 2010; **39** (suppl 1): i193–205.
- 78 Spears D. How much international variation in child height can sanitation explain? World Bank policy research working paper, no WPS 6351 2013. <http://go.worldbank.org/SZE5WUJB10> (accessed May 15, 2013).
- 79 Haider BA, Humayun Q, Bhutta ZA. Effect of administration of antihelminthics for soil transmitted helminths during pregnancy. *Cochrane Database Syst Rev* 2012; **2**: CD005547.
- 80 Taylor-Robinson D, Jones A, Garner P. Deworming drugs for treating soil-transmitted intestinal worms in children: effects on growth and school performance. *Cochrane Database Syst Rev* 2012; **4**: CD000371.
- 81 Gaffey MF, Wazny K, Bassani DG, Bhutta ZA. Dietary management of childhood diarrhea in low- and middle-income countries: a systematic review. *BMC Public Health* (in press).
- 82 Walker CLF, Black RE. Zinc for the treatment of diarrhoea: effect on diarrhoea morbidity, mortality and incidence of future episodes. *Int J Epidemiol* 2010; **39** (suppl 1): i63–69.
- 83 Garner P, Gulmezoglu AM. Drugs for preventing malaria in pregnant women. *Cochrane Database Syst Rev* 2009; **4**: CD000169.
- 84 Gamble C, Ekwuru JP, ter Kuile FO. Insecticide-treated nets for preventing malaria in pregnancy. *Cochrane Database Syst Rev* 2009; **2**: CD003755.
- 85 Meremikwu MM, Donegan S, Sinclair D, Esu E, Oranganje C. Intermittent preventive treatment for malaria in children living in areas with seasonal transmission. *Cochrane Database Syst Rev* 2012; **2**: CD003756.
- 86 Lengeler C. Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev* 2009; **2**: CD000363.
- 87 Dangour AD, Watson L, Cumming O, et al. Interventions to improve water quality and supply, sanitation and hygiene practices, and their effects on the nutritional status of children. *Cochrane Database Syst Rev* 2011; **3**: CD009382.
- 88 Awasthi S, Peto R, Read S, Richards SM, Pande V, Bundy D. Population deworming every 6 months with albendazole in 1 million pre-school children in north India: DEVTA, a cluster-randomised trial. *Lancet* 2013; published online March 14. [http://dx.doi.org/10.1016/S0140-6736\(12\)62126-6](http://dx.doi.org/10.1016/S0140-6736(12)62126-6).
- 89 WHO. Guideline update: technical aspects of the management of severe acute malnutrition in infants and children. Geneva: World Health Organization, 2013.
- 90 Collins S, Sadler K, Dent N, et al. Key issues in the success of community-based management of severe malnutrition. *Food Nutr Bull* 2006; **27**: S49–82.
- 91 Schofield C, Ashworth A. Why have mortality rates for severe malnutrition remained so high? *Bull World Health Organ* 1996; **74**: 223–29.
- 92 Chinkhumba J, Tomkins A, Banda T, Mkangama C, Fergusson P. The impact of HIV on mortality during in-patient rehabilitation of severely malnourished children in Malawi. *Trans R Soc Trop Med Hyg* 2008; **102**: 639–44.
- 93 Fergusson P, Chinkhumba J, Grijalva-Eternod C, Banda T, Mkangama C, Tomkins A. Nutritional recovery in HIV-infected and HIV-uninfected children with severe acute malnutrition. *Arch Dis Child* 2009; **94**: 512–16.
- 94 Hossain MM, Hassan MQ, Rahman MH, Kabir A, Hannan AH, Rahman A. Hospital management of severely malnourished children: comparison of locally adapted protocol with WHO protocol. *Indian Pediatr* 2009; **46**: 213–17.
- 95 Khanum S, Ashworth A, Huttly SR. Growth, morbidity, and mortality of children in Dhaka after treatment for severe malnutrition: a prospective study. *Am J Clin Nutr* 1998; **67**: 940–45.
- 96 Puaane T, Cuming K, Sanders D, Ashworth A. Why do some hospitals achieve better care of severely malnourished children than others? Five-year follow-up of rural hospitals in Eastern Cape, South Africa. *Health Policy Plan* 2008; **23**: 428–37.
- 97 Lenters LM, Wazny K, Webb P, Ahmed T, Bhutta ZA. Treatment of severe and moderate acute malnutrition in low- and middle-income settings: a systematic review, meta-analysis and delphi process. *BMC Public Health* (in press).
- 98 Trehan I, Goldbach HS, LaGrone LN, et al. Antibiotics as part of the management of severe acute malnutrition. *N Engl J Med* 2013; **368**: 425–35.
- 99 Collins S. Community-based therapeutic care—a new paradigm for selective feeding in nutritional crises. Humanitarian Policy Network paper 48. London: Overseas Development Institute, 2004.
- 100 Furber CM, McGowan L, Bower P, Kontopantelis E, Quenby S, Lavender T. Antenatal interventions for reducing weight in obese women for improving pregnancy outcome. *Cochrane Database Syst Rev* 2013; **1**: CD009334.
- 101 Amorin Adegboye AR, Linne YM, Lourenco PMC. Diet or exercise, or both, for weight reduction in women after childbirth. *Cochrane Database Syst Rev* 2012; **2**: CD005627.
- 102 Arenz S, Ruckerl R, Koletzko B, von Kries R. Breast-feeding and childhood obesity—a systematic review. *Int J Obes Relat Metab Disord* 2004; **28**: 1247–56.
- 103 Harder T, Bergmann R, Kallschnigg G, Plagemann A. Duration of breastfeeding and risk of overweight: a meta-analysis. *Am J Epidemiol* 2005; **162**: 397–403.
- 104 Owen CG, Martin RM, Whincup PH, Davey-Smith G, Gillman MW, Cook DG. The effect of breastfeeding on mean body mass index throughout life: a quantitative review of published and unpublished observational evidence. *Am J Clin Nutr* 2005; **82**: 1298–307.
- 105 Owen CG, Martin RM, Whincup PH, Smith GD, Cook DG. Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics* 2005; **115**: 1367–77.
- 106 Plagemann A, Harder T. Breast feeding and the risk of obesity and related metabolic diseases in the child. *Metab Syndr Relat Disord* 2005; **3**: 222–32.
- 107 Waters E, de Silva-Sanigorski A, Hall BJ, et al. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2011; **12**: CD001871.
- 108 Martin RM, Patel R, Kramer MS, et al. Effects of promoting longer-term and exclusive breastfeeding on adiposity and insulin-like growth factor-1 at age 11.5 years: a randomized trial. *JAMA* 2013; **309**: 1005–13.
- 109 Oude Luttikhuis H, Baur L, Jansen H, et al. Interventions for treating obesity in children. *Cochrane Database Syst Rev* 2009; **1**: CD001872.
- 110 Horton S, Alderman H, Rivera J. Hunger and malnutrition. Copenhagen consensus 2008: malnutrition and hunger. <http://thousanddays.org/wp-content/uploads/2011/05/Copenhagen-Consensus-2008-summary.pdf> (accessed March 15, 2013).
- 111 WHO, FAO. Guidelines on food fortification with micronutrients. Geneva: World Health Organization; Food and Agricultural Organization, 2006.
- 112 Mayer JE, Pfeiffer WH, Beyer P. Biofortified crops to alleviate micronutrient malnutrition. *Curr Opin Plant Biol* 2008; **11**: 166–70.
- 113 Eichler K, Wieser S, Ruthemann I, Brugger U. Effects of micronutrient fortified milk and cereal food for infants and children: a systematic review. *BMC Public Health* 2012; **12**: 506.
- 114 Gera T, Sachdev HS, Boy E. Effect of iron-fortified foods on hematologic and biological outcomes: systematic review of randomized controlled trials. *Am J Clin Nutr* 2012; **96**: 309–24.
- 115 Natri AM, Salo P, Vikstedt T, et al. Bread fortified with cholecalciferol increases the serum 25-hydroxyvitamin D concentration in women as effectively as a cholecalciferol supplement. *J Nutr* 2006; **136**: 123–27.
- 116 Diaz-Gomez NM, Domenech E, Barroso F, Castells S, Cortabarra C, Jimenez A. The effect of zinc supplementation on linear growth, body composition, and growth factors in preterm infants. *Pediatrics* 2003; **111**: 1002–09.
- 117 Harvey PWJ, Dary O. Governments and academic institutions play vital roles in food fortification: iron as an example. *Public Health Nutr* 2012; **1**: 1–5.
- 118 Mannar MG, van Ameringen M. Role of public-private partnership in micronutrient food fortification. *Food Nutr Bull* 2003; **24** (suppl 4): S151–54.
- 119 Gillespie S, Haddad L, Mannar V, Menon P, Nisbett N, and the Maternal and Child Nutrition Study Group. The politics of reducing malnutrition: building commitment and accelerating progress. *Lancet* 2013; published online June 6. [http://dx.doi.org/10.1016/S0140-6736\(13\)60842-9](http://dx.doi.org/10.1016/S0140-6736(13)60842-9).

- 120 Bassani DG, Arora P, Wazny K, Gaffey MF, Lenters L, Bhutta ZA. Financial incentives and coverage of child health interventions: a systematic review and meta-analysis. *BMC Public Health* (in press).
- 121 Haines A, Sanders D, Lehmann U, et al. Achieving child survival goals: potential contribution of community health workers. *Lancet* 2007; **369**: 2121–31.
- 122 Bhutta ZA, Das JK, Walker N, et al, for *The Lancet* Diarrhoea and Pneumonia Interventions Study Group. Interventions to address deaths from childhood pneumonia and diarrhoea equitably: what works and at what cost? *Lancet* 2013; **381**: 1417–29.
- 123 Lassi ZS, Haider BA, Bhutta ZA. Community-based intervention packages for reducing maternal and neonatal morbidity and mortality and improving neonatal outcomes. *Cochrane Database Syst Rev* 2010; **11**: CD007754.
- 124 Lewin S, Munabi-Babigumira S, Glenton C, et al. Lay health workers in primary and community health care for maternal and child health and the management of infectious diseases. *Cochrane Database Syst Rev* 2010; **3**: CD004015.
- 125 GHWA. Global experience of community health workers for delivery of health related millennium development goals: a systematic review, country case studies, and recommendations for integration into national health systems. Geneva: Global Health Workforce Alliance, 2010.
- 126 Gove S. Integrated management of childhood illness by outpatient health workers: technical basis and overview. The WHO Working Group on Guidelines for Integrated Management of the Sick Child. *Bull World Health Organ* 1997; **75** (suppl 1): 7–24.
- 127 Ahmed HM, Mitchell M, Hedt B. National implementation of Integrated Management of Childhood Illness (IMCI): policy constraints and strategies. *Health Policy* 2010; **96**: 128–33.
- 128 Schellenberg JRM, Adam T, Mshinda H, et al. Effectiveness and cost of facility-based Integrated Management of Childhood Illness (IMCI) in Tanzania. *Lancet* 2004; **364**: 1583–94.
- 129 Arifeen SE, Hoque DM, Akter T, et al. Effect of the Integrated Management of Childhood Illness strategy on childhood mortality and nutrition in a rural area in Bangladesh: a cluster randomised trial. *Lancet* 2009; **374**: 393–403.
- 130 Kristjansson B, Petticrew M, MacDonald B, et al. School feeding for improving the physical and psychosocial health of disadvantaged students. *Cochrane Database Syst Rev* 2009; **1**: CD004676.
- 131 Doherty T, Chopra M, Tomlinson M, Oliphant N, Nsibande D, Mason J. Moving from vertical to integrated child health programmes: experiences from a multi-country assessment of the Child Health Days approach in Africa. *Trop Med Int Health* 2010; **15**: 296–305.
- 132 UNICEF. Child health days 1999–2009: key achievements and the way forward. A report prepared for the UNICEF Joint Working Group on Child Health Days. New York, NY: UNICEF, 2011.
- 133 The Sphere Project. Sphere handbook. 2011. <http://www.spherehandbook.org/en/how-to-use-this-chapter-3/> (accessed May 15, 2013).
- 134 Young H, Borrel A, Holland D, Salama P. Public nutrition in complex emergencies. *Lancet* 2004; **364**: 1899–909.
- 135 World Food Programme. Nutrition in emergencies: WFP experiences and challenges. *Food Nutr Bull* 2006; **27**: 57–66.
- 136 The Johns Hopkins University's Bloomberg School of Public Health and International Committee of the Red Cross. Public health guide in emergencies, 2nd edn. Geneva: ICRC, 2008.
- 137 Global Nutrition Cluster of the Inter Agency Standing Committee. A toolkit for addressing nutrition in emergency situations. New York, NY: United Nations International Children's Fund, 2008.
- 138 Bailey S, Hedlund K. The impact of cash transfers on nutrition in emergency and transitional contexts: a review of the evidence. HPG Synthesis Paper. London: Overseas Development Institute, 2012.
- 139 Langendorf C, Roederer T. Evaluation de différentes stratégies de distributions préventives de la malnutrition au Niger. 2012 District de Madarounfa Région de Maradi Août 2011–Novembre 2012. Rapport intermédiaire. Paris: Médecins Sans Frontières, 2012.
- 140 Grijalva-Eternod CS, Wells JCK, Cortina-Borja M, et al. The double burden of obesity and malnutrition in a protracted emergency setting: a cross-sectional study of western Sahara refugees. *PLoS Med* 2012; **9**: e1001320.
- 141 IFRC. Focus on hunger and malnutrition. World Disasters Report 2011. Geneva: International Federation of Red Cross and Red Crescent Societies, 2011.
- 142 Hall A, Blankson B, Shoham J. The impact and effectiveness of emergency nutrition and Nutrition related interventions: a review of published evidence 2004–2010. Oxford: Emergency Nutrition Network, 2011.
- 143 UNHCR, WFP. Guidelines for selective feeding: the management of malnutrition in emergencies. In collaboration with the United Nations Standing Committee on Nutrition and the World Health Organization. Geneva: United Nations High Commission for Refugees, World Food Programme, 2011.
- 144 WHO, WFP, SCN, UNICEF. Community-Based Management of Severe Acute Malnutrition: A Joint Statement by the World Health Organization, the World Food Programme, the United Nations System Standing Committee on Nutrition and the United Nations Children's Fund. Geneva: World Health Organization, World Food Programme, United Nations System Standing Committee on Nutrition, United Nations Children's Fund, 2007.
- 145 Jones G, Steketee RW, Black RE, Bhutta ZA, Morris SS, and the Bellagio Child Survival Study Group. How many child deaths can we prevent this year? *Lancet* 2003; **362**: 65–71.
- 146 Dherani M, Pope D, Mascarenhas M, Smith KR, Weber M, Bruce NG. Indoor air pollution from unprocessed solid fuel use and pneumonia risk in under-5 children: systematic review and meta-analysis. *Bull World Health Organ* 2008; **86**: 390–98.
- 147 Pope DP, Mishra V, Thompson L, et al. Risk of low birth weight and stillbirth associated with indoor air pollution from solid fuel use in developing countries. *Epidemiol Rev* 2010; **32**: 70–81.
- 148 Bruce N, Dherani M, Das J, et al. Control of household air pollution for child survival: estimates for intervention impacts. *BMC Public Health* (in press).
- 149 Smith KR, McCracken JP, Weber MW, et al. Effect of reduction in household air pollution on childhood pneumonia in Guatemala (RESPIRE): a randomised controlled trial. *Lancet* 2011; **378**: 1717–26.
- 150 Thompson LM, Bruce N, Eskenazi B, Diaz A, Pope D, Smith KR. Impact of reduced maternal exposures to wood smoke from an introduced chimney stove on newborn birth weight in rural Guatemala. *Environ Health Perspect* 2011; **119**: 1489–94.
- 151 Wei SQ, Qi HP, Luo ZC, Fraser WD. Maternal vitamin D status and adverse pregnancy outcomes: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med* 2013; published online Feb 11. DOI:10.31009/14767058.2013.765849.
- 152 Lawlor DA, Wills AK, Fraser A, Sayers A, Fraser WD, Tobias JH. Association of maternal vitamin D status during pregnancy with bone-mineral content in offspring: a prospective cohort study. *Lancet* 2013; published online March 19. [http://doi.dx.org/10.1016/S0140-6736\(12\)62203-X](http://doi.dx.org/10.1016/S0140-6736(12)62203-X).
- 153 De-Regil LM, Palacios C, Ansary A, Kulier R, Pena-Rosas JP. Vitamin D supplementation for women during pregnancy. *Cochrane Database Syst Rev* 2012; **2**: CD008873.
- 154 Mori R, Ota E, Middleton P, Tobe-Gai R, Mahomed K, Bhutta ZA. Zinc supplementation for improving pregnancy and infant outcome. *Cochrane Database Syst Rev* 2012; **7**: CD000230.
- 155 Makrides M, Duley L, Olsen SF. Marine oil, and other prostaglandin precursor, supplementation for pregnancy uncomplicated by pre-eclampsia or intrauterine growth restriction. *Cochrane Database Syst Rev* 2007; **3**: CD003402.
- 156 Salvig JD, Lamont RF. Evidence regarding an effect of marine n-3 fatty acids on preterm birth: a systematic review and meta-analysis. *Acta Obstet Gynecol Scand* 2011; **90**: 825–38.
- 157 Dziechciarz P, Horvath A, Szajewska H. Effects of n-3 long-chain polyunsaturated fatty acid supplementation during pregnancy and/or lactation on neurodevelopment and visual function in children: a systematic review of randomized controlled trials. *J Am Coll Nutr* 2010; **29**: 443–54.
- 158 Klemens CM, Berman DR, Mozurkewich EL. The effect of perinatal omega-3 fatty acid supplementation on inflammatory markers and allergic diseases: a systematic review. *BJOG* 2011; **118**: 916–25.
- 159 Rodriguez G, Iglesia I, Bel-Serrat S, Moreno LA. Effect of n-3 long chain polyunsaturated fatty acids during the perinatal period on later body composition. *Br J Nutr* 2012; **107**: S117–28.
- 160 Szajewska H, Horvath A, Koletzko B. Effect of n-3 long-chain polyunsaturated fatty acid supplementation of women with low-risk pregnancies on pregnancy outcomes and growth measures at birth: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2006; **83**: 1337–44.

- 161 Ryan AS, Astwood JD, Gautier S, Kuratko CN, Nelson EB, Salem N Jr. Effects of long-chain polyunsaturated fatty acid supplementation on neurodevelopment in childhood: a review of human studies. *Prostaglandins Leukot Essent Fatty Acids* 2010; **82**: 305–14.
- 162 Muhlhäusler BS, Gibson RA, Makrides M. Effect of long-chain polyunsaturated fatty acid supplementation during pregnancy or lactation on infant and child body composition: a systematic review. *Am J Clin Nutr* 2010; **92**: 857–63.
- 163 Michaelsen KF, Dewey KG, Perez-Exposito AB, Nurhasan M, Lauritzen L, Roos N. Food sources and intake of n-6 and n-3 fatty acids in low-income countries with emphasis on infants, young children (6–24 months), and pregnant and lactating women. *Matern Child Nutr* 2011; **7**: 124–40.
- 164 Prentice AM, van der Merwe L. Impact of fatty acid status on immune function of children in low-income countries. *Matern Child Nutr* 2011; **7**: 89–98.
- 165 Rahman A, Patel V, Maselko J, Kirkwood B. The neglected 'm' in MCH programmes—why mental health of mothers is important for child nutrition. *Trop Med Int Health* 2008; **13**: 579–83.
- 166 Satyanarayana VA, Lukose A, Srinivasan K. Maternal mental health in pregnancy and child behavior. *Indian J Psychiatr* 2011; **53**: 351.
- 167 Austin MP, Priest SR, Sullivan EA. Antenatal psychosocial assessment for reducing perinatal mental health morbidity. *Cochrane Database Syst Rev* 2008; **4**: CD005124.
- 168 Diego MA, Field T, Hernandez-Reif M, Deeds O, Ascencio A, Begert G. Preterm infant massage elicits consistent increases in vagal activity and gastric motility that are associated with greater weight gain. *Acta Paediatr* 2007; **96**: 1588–91.
- 169 Vickers A, Ohlsson A, Lacy JB, Horsley A. Massage for promoting growth and development of preterm and/or low birth-weight infants. *Cochrane Database Syst Rev* 2009; **2**: CD000390.
- 170 Field T, Diego M, Hernandez-Reif M. Preterm infant massage therapy research: a review. *Infant Behav Dev* 2011; **33**: 115–24.
- 171 Lerch C, Meissner T. Interventions for the prevention of nutritional rickets in term born children. *Cochrane Database Syst Rev* 2007; **4**: CD006164.
- 172 Haider BA, Lassi ZS, Ahmed A, Bhutta ZA. Zinc supplementation as an adjunct to antibiotics in the treatment of pneumonia in children 2 to 59 months of age. *Cochrane Database Syst Rev* 2011; **10**: CD007368.
- 173 Basnet S, Shrestha PS, Sharma A, et al. A randomized controlled trial of zinc as adjuvant therapy for severe pneumonia in young children. *Pediatrics* 2012; **129**: 701–08.
- 174 Das RR, Singh M, Shafiq N. Short-term therapeutic role of zinc in children <5 years of age hospitalized for severe acute lower respiratory tract infection. *Paediatr Respir Rev* 2012; **13**: 184–91.
- 175 Shah GS, Dutta AK, Shah D, Mishra OP. Role of zinc in severe pneumonia: a randomized double blind placebo controlled study. *Ital J Pediatr* 2012; **38**: 36.
- 176 Srinivasan MG, Ndeezzi G, Mboijana CK, et al. Zinc adjunct therapy reduces case fatality in severe childhood pneumonia: a randomized double blind placebo-controlled trial. *BMC Med* 2012; **10**: 14.
- 177 Bhatnagar S, Wadhwa N, Aneja S, et al. Zinc as adjunct treatment in infants aged between 7 and 120 days with probable serious bacterial infection: a randomised, double-blind, placebo-controlled trial. *Lancet* 2012; **379**: 2072–78.
- 178 Mehta K, Bhatta NK, Majhi S, Shrivastava MK, Singh RR. Role of zinc in neonatal sepsis: a double blinded, randomized, placebo controlled trial. *Indian Pediatr* 2013; **50**: 390–93.
- 179 Phuka J, Maleta K, Thakwalakwa C. Complementary feeding with fortified spread and incidence of severe stunting in 6–8 month-old rural Malawians. *Arch Pediatr Adolesc Med* 2008; **162**: 619–26.
- 180 Phuka J, Maleta K, Thakwalakwa C, et al. Postintervention growth of Malawian children who received 12-mo dietary complementation with a lipid-based nutrient supplement or maize-soy flour. *Am J Clin Nutr* 2009; **89**: 382–90.
- 181 Adu-Afarwuah S, Lartey A, Brown KH, Zlotkin S, Briend A, Dewey KG. Randomized comparison of 3 types of micronutrient supplements for home fortification of complementary foods in Ghana: effects on growth and motor development. *Am J Clin Nutr* 2007; **86**: 412–20.
- 182 Adu-Afarwuah S, Lartey A, Brown KH, Zlotkin S, Briend A, Dewey KG. Home fortification of complementary foods with micronutrient supplements is well accepted and has positive effects on infant iron status in Ghana. *Am J Clin Nutr* 2008; **87**: 929–38.
- 183 Walker N, Friberg I. Overview of the Lives Saved Tool (LiST). *BMC Public Health* (in press).
- 184 Darmstadt G, Walker N, Lawn JE, Bhutta ZA, Haws RA, Cousens S. Saving newborn lives in Asia and Africa: cost and impact of phased scale-up of interventions within the continuum of care. *Health Policy Plann* 2008; **23**: 101–17.
- 185 Garnett GP, Cousens S, Hallett TB, Steketee, R, Walker N. Mathematical models in the evaluation of health programmes. *Lancet* 2011; **378**: 515–25.
- 186 Sachdev HPS, Hall A, Walker N, eds. Development and use of the Lives Saved Tool (LiST): a model to estimate the impact of scaling up proven interventions on maternal, neonatal and child mortality. *Int J Epidemiol* 2010; **39** (suppl 1): i1–205.
- 187 Fox M, Marterell R, van den Broek N, Walker N, eds. Technical inputs, enhancements and applications of the Lives Saved Tool (LiST). *BMC Public Health* 2011; **11** (suppl 3): S22.
- 188 Walker N, ed. Updates of assumptions and methods for the Lives Saved Tool (LiST). *BMC Public Health* (in press).
- 189 WHO, Global Health Observatory Data Repository. Life expectancy at birth, low income countries, both sexes combined, 2011. <http://apps.who.int/gho/data/view.main.700?lang=en> (accessed April 8, 2013).
- 190 Amouzou A, Richard SA, Friberg IK, et al. How well does LiST capture mortality by wealth quintile? A comparison of measured versus modelled mortality rates among children under-five in Bangladesh. *Int J Epidemiol* 2010; **39** (suppl 1): i186–92.
- 191 National Institute of Population Studies. Pakistan Demographic and Health Survey 2006–07. Islamabad, Pakistan. <http://www.measuredhs.com/pubs/pdf/FR200/FR200.pdf> (accessed on March 2, 2013).
- 192 Bhutta ZA, Hafeez A, Rizvi, et al. Reproductive, maternal, newborn, and child in Pakistan: challenges and opportunities. *Lancet* 2013; published online May 17 [http://dx.doi.org/10.1016/S0140-6736\(12\)61999-0](http://dx.doi.org/10.1016/S0140-6736(12)61999-0).
- 193 Futures Institute. UN integrated costing tool. <http://www.who.int/workforcealliance/knowledge/toolkit/10/en/> (accessed Dec 7, 2012).
- 194 Horton MS, McDonald C, Mahal A. Scaling-up nutrition: what will it cost? Washington DC: World Bank Directions in Development, 2009.
- 195 WHO. Health systems and information. http://www.who.int/healthinfo/global_burden_disease/definition_regions/en/ (accessed April 20, 2013).
- 196 Winkvist A, Rasmussen KM, Habicht JP. A new definition of maternal depletion syndrome. *Am J Public Health* 1992; **82**: 691–94.
- 197 Stevens GA, Finucane MM, Paciorek CJ, et al. Trends in mild, moderate, and severe stunting and underweight, and progress towards MDG 1 in 141 developing countries: a systematic analysis of population representative data. *Lancet* 2012; **380**: 824–34.
- 198 UNICEF. Improving child nutrition: the increasing imperative for global progress. New York, NY: United Nations International Children's Fund, 2013.
- 199 Victora CG, Barros AJ, Axelson H, et al. How changes in coverage affect equity in maternal and child health interventions in 35 Countdown to 2015 countries: an analysis of national surveys. *Lancet* 2012; **380**: 1149–56.
- 200 IHME. Financing global health 2012: the end of the golden age? Seattle, WA: Institute for Health Metrics and Evaluation, 2012.
- 201 Hsu J, Pitt C, Greco G, Berman P, Mills A. Countdown to 2015: changes in official development assistance to maternal, newborn, and child health in 2009–10, and assessment of progress since 2003. *Lancet* 2012; **380**: 1157–68.
- 202 Adair L, Fall CHD, Osmond C, et al, for the COHORTS group. Associations of linear growth and relative weight gain during early life with adult health and human capital in countries of low and middle income: findings from five birth cohort studies. *Lancet* 2013; published online March 28. [http://dx.doi.org/10.1016/S1404-6736\(13\)60103-8](http://dx.doi.org/10.1016/S1404-6736(13)60103-8).
- 203 Bhutta ZA, Chopra M. The Countdown for 2015: what lies ahead? *Lancet* 2012; **380**: 1125–27.